

Pseudosymmetric braidings, twines and twisted algebras

Florin Panaite*

Institute of Mathematics of the Romanian Academy
PO-Box 1-764, RO-014700 Bucharest, Romania
e-mail: Florin.Panaite@imar.ro

Mihai D. Staic†

Department of Mathematics, Indiana University,
Rawles Hall, Bloomington, IN 47405, USA
e-mail: mstaic@indiana.edu

Freddy Van Oystaeyen

Department of Mathematics and Computer Science
University of Antwerp, Middelheimlaan 1
B-2020 Antwerp, Belgium
e-mail: Francine.Schoeters@ua.ac.be

Abstract

A *laycle* is the categorical analogue of a lazy cocycle. *Twines* (as introduced by Bruguières) and *strong twines* (as introduced by the authors) are laycles satisfying some extra conditions. If c is a braiding, the double braiding c^2 is always a twine; we prove that it is a strong twine if and only if c satisfies a sort of modified braid relation (we call such c *pseudosymmetric*, as any symmetric braiding satisfies this relation). It is known that symmetric Yetter-Drinfeld categories are trivial; we prove that the Yetter-Drinfeld category ${}_H\mathcal{YD}^H$ over a Hopf algebra H is pseudosymmetric if and only if H is commutative and cocommutative. We introduce as well the Hopf algebraic counterpart of pseudosymmetric braidings under the name *pseudotriangular structures* and prove that all quasitriangular structures on the 2^{n+1} -dimensional pointed Hopf algebras $E(n)$ are pseudotriangular. We observe that a laycle on a monoidal category induces a so-called *pseudotwistor* on every algebra in the category, and we obtain some general results (and give some examples) concerning pseudotwistors, inspired by properties of laycles and twines.

*Research carried out while the first author was visiting the University of Antwerp, supported by a postdoctoral fellowship offered by FWO (Flemish Scientific Research Foundation). This author was also partially supported by the programme CEEX of the Romanian Ministry of Education and Research, contract nr. 2-CEX06-11-20/2006.

†Permanent address: Institute of Mathematics of the Romanian Academy, PO-Box 1-764, RO-014700 Bucharest, Romania.

Introduction

The notion of *symmetric category* is a classical concept in category theory. It consists of a monoidal category $\mathcal{C}$ equipped with a family of natural isomorphisms $c_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ satisfying natural “bilinearity” conditions together with the symmetry relation $c_{Y,X} \circ c_{X,Y} = id_{X \otimes Y}$, for all $X, Y \in \mathcal{C}$. In 1985 Joyal and Street were led by natural considerations to drop this symmetry condition from the axioms, thus arriving at the concept of *braiding*, which afterwards became of central importance for the then emerging theory of quantum groups; for instance, if (H, R) is a quasitriangular Hopf algebra as defined by Drinfeld, then the monoidal category ${}_H\mathcal{M}$ of left H -modules acquires a braiding defined by R , which is symmetric if and only if R is triangular, i.e. $R_{21}R = 1 \otimes 1$.

There exist many examples of symmetric braidings, as well as many examples of braidings which are not symmetric. Although some of the most basic examples of monoidal categories (such as the category of vector spaces) are symmetric, the symmetry condition is a rather restrictive requirement, a claim which is probably best illustrated by the following result of Pareigis (cf. [30]): if H is a Hopf algebra, then the Yetter-Drinfeld category ${}_H\mathcal{YD}^H$ is symmetric if and only if H is trivial (i.e. $H = k$). Thus, the most basic examples of braided categories arising in Hopf algebra theory are virtually never symmetric.

It appears thus natural to look for braidings satisfying some generalized (or weakened) symmetry conditions. In a recent paper [15], Etingof and Gelaki proposed the concept of *quasisymmetric braiding*, as being a braiding with the property that $c_{Y,X} \circ c_{X,Y} = id_{X \otimes Y}$ for all X, Y *simple* objects in the category, and classified quasisymmetric braided categories of exponential growth, generalizing Deligne’s classification of symmetric categories of exponential growth. On the other hand, at the Hopf algebraic level, Liu and Zhu proposed in [23] the concept of *almost-triangular* Hopf algebra, as being a quasitriangular Hopf algebra (H, R) such that $R_{21}R$ is central in $H \otimes H$ (obviously, this concept generalizes the one of triangular Hopf algebra, but it is not clear whether it has a categorical counterpart).

The original aim of the present paper was to continue the study of some categorical concepts recently introduced in [34], [5], [29] under the names *pure-braided structure*, *twine* and *strong twine*. We recall from [5] that a twine on a monoidal category $\mathcal{C}$ is a family of natural isomorphisms $D_{X,Y} : X \otimes Y \rightarrow X \otimes Y$ in $\mathcal{C}$ satisfying a certain list of axioms chosen in such a way that, if c is a braiding on $\mathcal{C}$, then the so-called *double braiding* c^2 defined by $c_{X,Y}^2 = c_{Y,X} \circ c_{X,Y}$ is a twine (by [29], the concept of twine is equivalent to the concept of pure-braided structure introduced in [34]). Moreover, twines are related to the pure braid groups in the same way in which braidings are related to the braid groups. A strong twine, as defined in [29], is also a family of natural isomorphisms $D_{X,Y} : X \otimes Y \rightarrow X \otimes Y$ in $\mathcal{C}$ satisfying a list of (easier looking) axioms, which imply the axioms of a twine. A double braiding c^2 is *not* always a strong twine, so we were led naturally to ask for what kind of braidings c is c^2 a strong twine. The answer is that this happens if and only if c satisfies the following condition:

$$(c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1}) \circ (c_{X,Y} \otimes id_Z) = (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y) \circ (id_X \otimes c_{Y,Z})$$

for all $X, Y, Z \in \mathcal{C}$. This is a sort of modified braid relation, and it is obvious that if c is a symmetry then this condition becomes exactly the braid relation satisfied by any braiding; thus, any symmetric braiding satisfies the above relation, so what we obtained is a generalized symmetry condition. A braiding satisfying the above modified braid relation will be called *pseudosymmetric*. It should be emphasized that, although we arrived at this concept in an indirect way (via double braidings and strong twines), the pseudosymmetry relation does not

depend on these concepts and could have been introduced directly. Anyway, this concept is supported and further justified by our main result: if H is a Hopf algebra (with bijective antipode) then the canonical braiding of the Yetter-Drinfeld category ${}_H\mathcal{YD}^H$ is pseudosymmetric if and only if H is commutative and cocommutative. In view of Pareigis' result mentioned above, this shows that pseudosymmetries are far more numerous than symmetries; and in the opposite direction, it shows that *not* every braiding is pseudosymmetric (this was not so obvious a priori). Note also that, incidentally, our theorem provides a characterization of commutative and cocommutative Hopf algebras solely in terms of their Yetter-Drinfeld categories.

We introduce the Hopf algebraic counterpart of pseudosymmetric braidings, under the name *pseudotriangular structure*, as being a quasitriangular structure R on a Hopf algebra H satisfying the modified quantum Yang-Baxter equation $R_{12}R_{31}^{-1}R_{23} = R_{23}R_{31}^{-1}R_{12}$ (from which it is visible that triangular implies pseudotriangular) or equivalently the element $F = R_{21}R$ satisfies the condition $F_{12}F_{23} = F_{23}F_{12}$, which shows immediately that almost-triangular implies pseudotriangular. We analyze in detail a class of quasitriangular Hopf algebras, namely the 2^{n+1} -dimensional pointed Hopf algebras $E(n)$ whose quasitriangular structures and cleft extensions have been classified in [27] and [28]: we prove that all quasitriangular structures of $E(n)$ (which are in bijection with $n \times n$ matrices) are pseudotriangular, and the only almost-triangular structures of $E(n)$ are the triangular ones (which are in bijection with symmetric $n \times n$ matrices); in particular, this shows that pseudotriangular does *not* imply almost-triangular.

Apart from leading us to consider a certain class of braidings (the pseudosymmetric ones), the study of twines led us also to consider certain classes of *pseudotwistors*, as introduced in [24]. In order to explain this, we need to introduce first some terminology. A basic object we use all over the paper is a monoidal structure of the identity functor on a monoidal category (for instance, this is part of the axioms for twines and strong twines). We needed to have a name for such an object, and in order to choose it we relied on the fact that these objects are the categorical analogues of *lazy cocycles*, a concept recently introduced in Hopf algebra theory and studied in a series of papers ([1], [7], [8], [9], [10], [33]). Thus, we have chosen the name **laycle**, as derived from **lazy cocycle**. These laycles have some properties similar to those of lazy cocycles, for instance they act by conjugation on braidings and it is possible to define for them an analogue of the Hopf lazy cohomology.

The concept of pseudotwistor (with particular cases called *twistor* and *braided twistor*) was introduced in [24] as an abstract and axiomatic device for “twisting” the multiplication of an algebra in a monoidal category in order to obtain a new algebra structure (on the same object). More precisely, if (A, μ, u) is an algebra in a monoidal category $\mathcal{C}$, a pseudotwistor for A is a morphism $T : A \otimes A \rightarrow A \otimes A$ in $\mathcal{C}$, for which there exist two morphisms $\tilde{T}_1, \tilde{T}_2 : A \otimes A \otimes A \rightarrow A \otimes A \otimes A$ in $\mathcal{C}$, called the companions of T , satisfying a list of axioms ensuring that $(A, \mu \circ T, u)$ is also an algebra in $\mathcal{C}$. Examples of pseudotwistors are abundant, cf. [24]. For instance, if c is a braiding on $\mathcal{C}$, then $c_{A,A}^2$ is a pseudotwistor for every algebra A in $\mathcal{C}$. Since a double braiding is in particular a twine, this raises the natural question whether any twine induces a pseudotwistor on every algebra in the category. It turns out that something more general holds, namely that any laycle has this property. This seems to show that pseudotwistors are “local” versions of laycles (in the same sense in which twisting maps are “local” versions of braidings, see [19] for the meaning of these concepts and references), but this is not quite true, because for instance a composition of laycles is a laycle while a composition of pseudotwistors is not in general a pseudotwistor. We introduce thus the concept of *strong pseudotwistor*, as a better candidate for being a local version of laycles (for instance, a composition of a strong pseudotwistor with itself is again a strong pseudotwistor). We also introduce a sort of local version of twines, under

the name *pure pseudotwistor*, as being a pseudotwistor whose companions satisfy the condition $(\tilde{T}_2 \otimes id) \circ (id \otimes \tilde{T}_1) = (id \otimes \tilde{T}_1) \circ (\tilde{T}_2 \otimes id)$. Quite interestingly, it turns out that virtually all the concrete examples of pseudotwistors we are aware of are pure.

What we discussed above are basically facts about pseudotwistors inspired by properties of laycles and twines. In the last section of the paper we complete the picture of the interplay between laycles and twines, on the one hand, and pseudotwistors, on the other hand, by presenting a result in the opposite direction. Namely, inspired by a result in [24] concerning pseudotwistors and twisting maps, we prove that, if $\mathcal{C}$ is a monoidal category, T a laycle and d a braiding on $\mathcal{C}$ related in a certain way, then the families $d'_{X,Y} = d_{X,Y} \circ T_{X,Y}$ and $d''_{X,Y} = T_{Y,X} \circ d_{X,Y}$ are also braidings on $\mathcal{C}$. We prove also a sort of converse result, leading thus to a characterization of generalized double braidings (i.e. twines of the type $c'_{Y,X} \circ c_{X,Y}$, with c, c' braidings).

1 Preliminaries

In this section we recall basic definitions and results and we fix notation to be used throughout the paper. All algebras, linear spaces, etc, will be over a base field k ; unadorned $\otimes$ means $\otimes_k$. All monoidal categories are assumed to be strict, with unit denoted by I . For a Hopf algebra H with comultiplication Δ we denote $\Delta(h) = h_1 \otimes h_2$, for all $h \in H$. Unless otherwise stated, H will denote a Hopf algebra with bijective antipode S . For terminology concerning Hopf algebras and monoidal categories we refer to [21], [26].

A linear map $\sigma : H \otimes H \rightarrow k$ is called a **left 2-cocycle** if it satisfies the condition

$$\sigma(a_1, b_1)\sigma(a_2 b_2, c) = \sigma(b_1, c_1)\sigma(a, b_2 c_2), \quad (1.1)$$

for all $a, b, c \in H$, and it is called a **right 2-cocycle** if it satisfies the condition

$$\sigma(a_1 b_1, c)\sigma(a_2, b_2) = \sigma(a, b_1 c_1)\sigma(b_2, c_2). \quad (1.2)$$

Given a linear map $\sigma : H \otimes H \rightarrow k$, define a product $\cdot_\sigma$ on H by $h \cdot_\sigma h' = \sigma(h_1, h'_1)h_2 h'_2$, for all $h, h' \in H$. Then $\cdot_\sigma$ is associative if and only if σ is a left 2-cocycle. If we define $\cdot_\sigma$ by $h \cdot_\sigma h' = h_1 h'_1 \sigma(h_2, h'_2)$, then $\cdot_\sigma$ is associative if and only if σ is a right 2-cocycle. In any of the two cases, σ is normalized (i.e. $\sigma(1, h) = \sigma(h, 1) = \varepsilon(h)$ for all $h \in H$) if and only if 1_H is the unit for $\cdot_\sigma$. If σ is a normalized left (respectively right) 2-cocycle, we denote the algebra $(H, \cdot_\sigma)$ by ${}_\sigma H$ (respectively H_σ). It is well-known that ${}_\sigma H$ (respectively H_σ) is a right (respectively left) H -comodule algebra via the comultiplication Δ of H . If $\sigma : H \otimes H \rightarrow k$ is normalized and convolution invertible, then σ is a left 2-cocycle if and only if σ^{-1} is a right 2-cocycle.

If $\gamma : H \rightarrow k$ is linear, normalized (i.e. $\gamma(1) = 1$) and convolution invertible, define

$$D^1(\gamma) : H \otimes H \rightarrow k, \quad D^1(\gamma)(h, h') = \gamma(h_1)\gamma(h'_1)\gamma^{-1}(h_2 h'_2), \quad \forall h, h' \in H.$$

Then $D^1(\gamma)$ is a normalized and convolution invertible left 2-cocycle.

We recall from [1] some facts about lazy cocycles and lazy cohomology. The set $Reg^1(H)$ (respectively $Reg^2(H)$) consisting of normalized and convolution invertible linear maps $\gamma : H \rightarrow k$ (respectively $\sigma : H \otimes H \rightarrow k$), is a group with respect to the convolution product. An element $\gamma \in Reg^1(H)$ is called **lazy** if $\gamma(h_1)h_2 = h_1\gamma(h_2)$, for all $h \in H$. The set of lazy elements of $Reg^1(H)$, denoted by $Reg_L^1(H)$, is a central subgroup of $Reg^1(H)$. An element $\sigma \in Reg^2(H)$ is called **lazy** if

$$\sigma(h_1, h'_1)h_2 h'_2 = h_1 h'_1 \sigma(h_2, h'_2), \quad \forall h, h' \in H. \quad (1.3)$$

The set of lazy elements of $\text{Reg}^2(H)$, denoted by $\text{Reg}_L^2(H)$, is a subgroup of $\text{Reg}^2(H)$. We denote by $Z^2(H)$ the set of left 2-cocycles on H and by $Z_L^2(H)$ the set $Z^2(H) \cap \text{Reg}_L^2(H)$ of normalized and convolution invertible lazy 2-cocycles. If $\sigma \in Z_L^2(H)$, then the algebras ${}_\sigma H$ and H_σ coincide and will be denoted by $H(\sigma)$; moreover, $H(\sigma)$ is an H -bicomodule algebra via Δ .

It is well-known that in general the set $Z^2(H)$ of left 2-cocycles is not closed under convolution. One of the main features of lazy 2-cocycles is that the set $Z_L^2(H)$ is closed under convolution, and that the convolution inverse of an element $\sigma \in Z_L^2(H)$ is again a lazy 2-cocycle, so $Z_L^2(H)$ is a group under convolution. In particular, a lazy 2-cocycle is also a right 2-cocycle.

Consider now the map $D^1 : \text{Reg}^1(H) \rightarrow \text{Reg}^2(H)$, $D^1(\gamma)(h, h') = \gamma(h_1)\gamma(h'_1)\gamma^{-1}(h_2h'_2)$, for all $h, h' \in H$. Then, by [1], the map D^1 induces a group morphism $\text{Reg}_L^1(H) \rightarrow Z_L^2(H)$, with image contained in the centre of $Z_L^2(H)$; denote by $B_L^2(H)$ this central subgroup $D^1(\text{Reg}_L^1(H))$ of $Z_L^2(H)$ (its elements are called **lazy 2-coboundaries**). Then define the **second lazy cohomology group** $H_L^2(H) = Z_L^2(H)/B_L^2(H)$.

Dually, an invertible element $T \in H \otimes H$ is called a **lazy twist** if

$$\begin{aligned} (\varepsilon \otimes \text{id})(T) &= 1 = (\text{id} \otimes \varepsilon)(T), \\ (\text{id} \otimes \Delta)(T)(1 \otimes T) &= (\Delta \otimes \text{id})(T)(T \otimes 1), \\ \Delta(h)T &= T\Delta(h), \quad \forall h \in H. \end{aligned}$$

As a consequence of these axioms we also have $(1 \otimes T)(\text{id} \otimes \Delta)(T) = (T \otimes 1)(\Delta \otimes \text{id})(T)$. One can define the analogues of $Z_L^2(H)$, $B_L^2(H)$ and $H_L^2(H)$ with lazy twists instead of lazy cocycles; these will be denoted respectively by $Z_{LT}^2(H)$, $B_{LT}^2(H)$ and $H_{LT}^2(H)$.

Remark 1.1 *If $\mathcal{C}$ is a monoidal category and $T_{X,Y} : X \otimes Y \rightarrow X \otimes Y$ is a family of natural isomorphisms in $\mathcal{C}$, the naturality of T implies (for all $X, Y, Z \in \mathcal{C}$):*

$$(T_{X,Y} \otimes \text{id}_Z) \circ T_{X \otimes Y, Z} = T_{X \otimes Y, Z} \circ (T_{X,Y} \otimes \text{id}_Z), \quad (1.4)$$

$$(\text{id}_X \otimes T_{Y,Z}) \circ T_{X, Y \otimes Z} = T_{X, Y \otimes Z} \circ (\text{id}_X \otimes T_{Y,Z}). \quad (1.5)$$

Definition 1.2 ([21]) *Let $\mathcal{C} = (\mathcal{C}, \otimes, I)$ and $\mathcal{D} = (\mathcal{D}, \otimes, I)$ be monoidal categories. A **monoidal functor** from $\mathcal{C}$ to $\mathcal{D}$ is a triple $(F, \varphi_0, \varphi_2)$ where $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor, φ_0 is an isomorphism in $\mathcal{D}$ from I to $F(I)$ and $\varphi_2(U, V) : F(U) \otimes F(V) \rightarrow F(U \otimes V)$ is a family of natural isomorphisms in $\mathcal{D}$ indexed by all couples (U, V) of objects in $\mathcal{C}$ such that, for all $U, V, W \in \mathcal{C}$:*

$$\begin{aligned} \varphi_2(U \otimes V, W) \circ (\varphi_2(U, V) \otimes \text{id}_{F(W)}) &= \varphi_2(U, V \otimes W) \circ (\text{id}_{F(U)} \otimes \varphi_2(V, W)), \\ \varphi_2(I, U) \circ (\varphi_0 \otimes \text{id}_{F(U)}) &= \text{id}_{F(U)}, \\ \varphi_2(U, I) \circ (\text{id}_{F(U)} \otimes \varphi_0) &= \text{id}_{F(U)}. \end{aligned}$$

Definition 1.3 ([20]) *Let $\mathcal{C}$ be a monoidal category. A **braiding** on $\mathcal{C}$ consists of a family of natural isomorphisms $c_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ in $\mathcal{C}$ such that, for all $X, Y, Z \in \mathcal{C}$:*

$$c_{X, Y \otimes Z} = (\text{id}_Y \otimes c_{X, Z}) \circ (c_{X, Y} \otimes \text{id}_Z), \quad (1.6)$$

$$c_{X \otimes Y, Z} = (c_{X, Z} \otimes \text{id}_Y) \circ (\text{id}_X \otimes c_{Y, Z}). \quad (1.7)$$

As consequences of the axioms we also have $c_{X, I} = c_{I, X} = \text{id}_X$ and the braid relation

$$\begin{aligned} (c_{Y, Z} \otimes \text{id}_X) \circ (\text{id}_Y \otimes c_{X, Z}) \circ (c_{X, Y} \otimes \text{id}_Z) \\ = (\text{id}_Z \otimes c_{X, Y}) \circ (c_{X, Z} \otimes \text{id}_Y) \circ (\text{id}_X \otimes c_{Y, Z}). \end{aligned} \quad (1.8)$$

*If moreover c satisfies $c_{Y, X} \circ c_{X, Y} = \text{id}_{X \otimes Y}$, for all $X, Y \in \mathcal{C}$, then c is called a **symmetry**.*

Definition 1.4 ([12]) Let H be a Hopf algebra. An invertible element $R \in H \otimes H$ is called a **quasitriangular structure** for H if

$$\begin{aligned}(\Delta \otimes id)(R) &= R_{13}R_{23}, \\(id \otimes \Delta)(R) &= R_{13}R_{12}, \\(\varepsilon \otimes id)(R) &= (id \otimes \varepsilon)(R) = 1, \\ \Delta^{cop}(h)R &= R\Delta(h), \quad \forall h \in H.\end{aligned}$$

If moreover R satisfies $R_{21}R = 1 \otimes 1$, then R is called **triangular**. If R is a quasitriangular (respectively triangular) structure for H , then the monoidal category ${}_H\mathcal{M}$ of left H -modules becomes braided (respectively symmetric), with braiding given by $c_{M,N} : M \otimes N \rightarrow N \otimes M$, $c_{M,N}(m \otimes n) = R^2 \cdot n \otimes R^1 \cdot m$, for all $M, N \in {}_H\mathcal{M}$, $m \in M$, $n \in N$.

Definition 1.5 ([34]) Let $\mathcal{C}$ be a monoidal category. A **pure-braided** structure on $\mathcal{C}$ consists of two families of natural isomorphisms $A_{U,V,W} : U \otimes V \otimes W \rightarrow U \otimes V \otimes W$ and $B_{U,V,W} : U \otimes V \otimes W \rightarrow U \otimes V \otimes W$ in $\mathcal{C}$ such that (for all $U, V, W, X \in \mathcal{C}$):

$$A_{U \otimes V, W, X} = A_{U, V \otimes W, X} \circ (id_U \otimes A_{V, W, X}), \quad (1.9)$$

$$A_{U, V, W \otimes X} = (A_{U, V, W} \otimes id_X) \circ A_{U, V \otimes W, X}, \quad (1.10)$$

$$B_{U \otimes V, W, X} = (id_U \otimes B_{V, W, X}) \circ B_{U, V \otimes W, X}, \quad (1.11)$$

$$B_{U, V, W \otimes X} = B_{U, V \otimes W, X} \circ (B_{U, V, W} \otimes id_X), \quad (1.12)$$

$$(A_{U, V, W} \otimes id_X) \circ (id_U \otimes B_{V, W, X}) = (id_U \otimes B_{V, W, X}) \circ (A_{U, V, W} \otimes id_X), \quad (1.13)$$

$$A_{U, I, V} = B_{U, I, V}. \quad (1.14)$$

A category equipped with a pure-braided structure is called a **pure-braided category**.

Definition 1.6 ([5]) Let $\mathcal{C}$ be a monoidal category. A **twine** on $\mathcal{C}$ is a family of natural isomorphisms $D_{X,Y} : X \otimes Y \rightarrow X \otimes Y$ in $\mathcal{C}$ satisfying the axioms (for all $X, Y, Z, W \in \mathcal{C}$):

$$D_{I, I} = id_I, \quad (1.15)$$

$$(D_{X,Y} \otimes id_Z) \circ D_{X \otimes Y, Z} = (id_X \otimes D_{Y,Z}) \circ D_{X, Y \otimes Z}, \quad (1.16)$$

$$\begin{aligned}(D_{X \otimes Y, Z} \otimes id_W) \circ (id_X \otimes D_{Y, Z}^{-1} \otimes id_W) \circ (id_X \otimes D_{Y, Z \otimes W}) \\ = (id_X \otimes D_{Y, Z \otimes W}) \circ (id_X \otimes D_{Y, Z}^{-1} \otimes id_W) \circ (D_{X \otimes Y, Z} \otimes id_W). \end{aligned} \quad (1.17)$$

A category equipped with a twine is called an **entwined category**. If $(\mathcal{C}, D)$ is entwined then we also have $D_{X, I} = D_{I, X} = id_X$, for all $X \in \mathcal{C}$.

By [29], these two concepts are equivalent in a certain (precise) sense.

Proposition 1.7 ([5]) Let $\mathcal{C}$ be a monoidal category and c, c' braidings on $\mathcal{C}$. Then the family $T_{X,Y} := c'_{Y,X} \circ c_{X,Y}$ is a twine, called a **generalized double braiding**; if $c = c'$ the family $c_{Y,X} \circ c_{X,Y}$ is called a **double braiding**.

Definition 1.8 ([29]) Let $\mathcal{C}$ be a monoidal category and $T_{X,Y} : X \otimes Y \rightarrow X \otimes Y$ a family of natural isomorphisms in $\mathcal{C}$. We say that T is a **strong twine** (or $(\mathcal{C}, T)$ is strongly entwined) if for all $X, Y, Z \in \mathcal{C}$ we have:

$$T_{I, I} = id_I, \quad (1.18)$$

$$(T_{X,Y} \otimes id_Z) \circ T_{X \otimes Y, Z} = (id_X \otimes T_{Y,Z}) \circ T_{X, Y \otimes Z}, \quad (1.19)$$

$$(T_{X,Y} \otimes id_Z) \circ (id_X \otimes T_{Y,Z}) = (id_X \otimes T_{Y,Z}) \circ (T_{X,Y} \otimes id_Z). \quad (1.20)$$

Proposition 1.9 ([29]) *If $(\mathcal{C}, T)$ is strongly entwined then $(\mathcal{C}, T)$ is entwined.*

Proposition 1.10 ([3], [4]) *Let A be an algebra with multiplication denoted by $\mu_A = \mu$ and let $T : A \otimes A \rightarrow A \otimes A$ be a linear map satisfying the following conditions: $T(1 \otimes a) = 1 \otimes a$, $T(a \otimes 1) = a \otimes 1$, for all $a \in A$, and*

$$\mu_{23} \circ T_{12} \circ T_{13} = T \circ \mu_{23} : A \otimes A \otimes A \rightarrow A \otimes A, \quad (1.21)$$

$$\mu_{12} \circ T_{23} \circ T_{13} = T \circ \mu_{12} : A \otimes A \otimes A \rightarrow A \otimes A, \quad (1.22)$$

$$T_{12} \circ T_{13} \circ T_{23} = T_{23} \circ T_{13} \circ T_{12} : A \otimes A \otimes A \rightarrow A \otimes A \otimes A, \quad (1.23)$$

with standard notation for μ_{ij} and T_{ij} . Then the map $\mu \circ T : A \otimes A \rightarrow A$ defines an associative algebra structure on A , with the same unit 1. The map T is called an **R-matrix** for A .

2 Laycles and quasi-braidings

Definition 2.1 *Let $\mathcal{C}$ be a monoidal category and $T_{X,Y} : X \otimes Y \rightarrow X \otimes Y$ a family of natural isomorphisms in $\mathcal{C}$. We say that T is a **laycle** if for all $X, Y, Z \in \mathcal{C}$ we have:*

$$T_{I,I} = id_I, \quad (2.1)$$

$$(T_{X,Y} \otimes id_Z) \circ T_{X \otimes Y, Z} = (id_X \otimes T_{Y,Z}) \circ T_{X, Y \otimes Z}. \quad (2.2)$$

A category equipped with a laycle is called a **laycled category**.

Remark 2.2 *It T is a laycle on $\mathcal{C}$ then we also have $T_{X,I} = T_{I,X} = id_X$, for all $X \in \mathcal{C}$. Also, it is clear that if $(\mathcal{C}, T)$ is entwined then $(\mathcal{C}, T)$ is laycled.*

Remark 2.3 *It is obvious that T is a laycle if and only if $(id_{\mathcal{C}}, id_I, \varphi_2(X, Y) := T_{X,Y})$ is a monoidal functor from $\mathcal{C}$ to itself. So, directly from the properties of monoidal functors, it follows that the composition of two laycles is a laycle and the inverse of a laycle is a laycle.*

Example 2.4 Let H be a Hopf algebra, $\sigma \in Reg_L^2(H)$ and $\mathcal{C} = \mathcal{M}^H$, the category of right H -comodules, with tensor product $(m \otimes n)_{(0)} \otimes (m \otimes n)_{(1)} = (m_{(0)} \otimes n_{(0)}) \otimes m_{(1)} n_{(1)}$. Define $T_{M,N}(m \otimes n) = m_{(0)} \otimes n_{(0)} \sigma(m_{(1)}, n_{(1)})$, for all $M, N \in \mathcal{M}^H$, $m \in M$, $n \in N$. Then σ is a lazy 2-cocycle on H if and only if T is a laycle on $\mathcal{M}^H$. Dually, if $F = F^1 \otimes F^2 \in H \otimes H$ is invertible and satisfies $(\varepsilon \otimes id)(F) = (id \otimes \varepsilon)(F) = 1$, consider the category ${}_H\mathcal{M}$ of left H -modules, with tensor product given by $h \cdot (m \otimes n) = h_1 \cdot m \otimes h_2 \cdot n$, for all $M, N \in {}_H\mathcal{M}$, $m \in M$, $n \in N$; define $T_{M,N}(m \otimes n) = F^1 \cdot m \otimes F^2 \cdot n$. Then F is a lazy twist if and only if T is a laycle on ${}_H\mathcal{M}$.

If T is a laycle on $\mathcal{C}$, we define the families $T_{X,Y,Z}^b, T_{X,Y,Z}^f : X \otimes Y \otimes Z \rightarrow X \otimes Y \otimes Z$ (notation as in [5]) of natural isomorphisms in $\mathcal{C}$ associated to it, by

$$T_{X,Y,Z}^b := (id_X \otimes T_{Y,Z}^{-1}) \circ T_{X \otimes Y, Z} = T_{X, Y \otimes Z} \circ (T_{X,Y}^{-1} \otimes id_Z), \quad (2.3)$$

$$T_{X,Y,Z}^f := T_{X \otimes Y, Z} \circ (id_X \otimes T_{Y,Z}^{-1}) = (T_{X,Y}^{-1} \otimes id_Z) \circ T_{X, Y \otimes Z}. \quad (2.4)$$

Proposition 2.5 *Let $\mathcal{C}$ be a monoidal category.*

(i) *If T is a laycle on $\mathcal{C}$ then for all $U, V, W \in \mathcal{C}$ we have*

$$T_{U \otimes V, W, X}^f = T_{U, V \otimes W, X}^f \circ (id_U \otimes T_{V, W, X}^f), \quad (2.5)$$

$$T_{U, V, W \otimes X}^f = (T_{U, V, W}^f \otimes id_X) \circ T_{U, V \otimes W, X}^f. \quad (2.6)$$

Conversely, if $A_{U, V, W} : U \otimes V \otimes W \rightarrow U \otimes V \otimes W$ is a family of natural isomorphisms such that (2.5) and (2.6) with A instead of T^f hold, then $T_{U, V} := A_{U, I, V}$ is a laycle on $\mathcal{C}$.

(ii) *If T is a laycle on $\mathcal{C}$ then for all $U, V, W \in \mathcal{C}$ we have*

$$T_{U \otimes V, W, X}^b = (id_U \otimes T_{V, W, X}^b) \circ T_{U, V \otimes W, X}^b, \quad (2.7)$$

$$T_{U, V, W \otimes X}^b = T_{U, V \otimes W, X}^b \circ (T_{U, V, W}^b \otimes id_X). \quad (2.8)$$

Conversely, if $B_{U, V, W} : U \otimes V \otimes W \rightarrow U \otimes V \otimes W$ is a family of natural isomorphisms such that (2.7) and (2.8) with B instead of T^b hold, then $T_{U, V} := B_{U, I, V}$ is a laycle on $\mathcal{C}$.

Proof. We prove (i), while (ii) is similar and left to the reader. We compute:

$$\begin{aligned} T_{U \otimes V, W, X}^f &= T_{U \otimes V \otimes W, X} \circ (id_U \otimes id_V \otimes T_{W, X}^{-1}) \\ &= T_{U \otimes V \otimes W, X} \circ (id_U \otimes T_{V \otimes W, X}^{-1}) \\ &\quad \circ (id_U \otimes T_{V \otimes W, X}) \circ (id_U \otimes id_V \otimes T_{W, X}^{-1}) \\ &\stackrel{(2.2)}{=} T_{U, V \otimes W, X}^f \circ (id_U \otimes T_{V, W, X}^f), \end{aligned}$$

proving (2.5); the proof of (2.6) is similar and left to the reader.

Assume now that $A_{-, -, -}$ is a family of natural isomorphisms satisfying (2.5) and (2.6); then obviously the family $T_{U, V} = A_{U, I, V}$ consists also of natural isomorphisms. If in (2.5) we take $V = W = X = I$ we obtain $T_{U, I} = T_{U, I} \circ (id_U \otimes T_{I, I})$, hence $T_{I, I} = id_I$. If we take $W = I$ in (2.5) and $V = I$ in (2.6) we obtain

$$T_{U \otimes V, X} = A_{U, V, X} \circ (id_U \otimes T_{V, X}), \quad T_{U, W \otimes X} = (T_{U, W} \otimes id_X) \circ A_{U, W, X},$$

which together imply (2.2). $\square$

The categorical analogue of the operator D^1 from the Preliminaries looks as follows:

Proposition 2.6 ([5]) *Let $\mathcal{C}$ be a monoidal category and $R_X : X \rightarrow X$ a family of natural isomorphisms in $\mathcal{C}$ such that $R_I = id_I$. Then the family*

$$D^1(R)_{X, Y} := (R_X \otimes R_Y) \circ R_{X \otimes Y}^{-1} = R_{X \otimes Y}^{-1} \circ (R_X \otimes R_Y) \quad (2.9)$$

is a laycle on $\mathcal{C}$.

The next result (whose proof is straightforward and will be omitted) provides the categorical analogue of Hopf lazy cohomology:

Proposition 2.7 *Let $\mathcal{C}$ be a small monoidal category. Then:*

(i) *If we denote by $Reg_L^1(\mathcal{C})$ the set of families of natural isomorphisms $R_X : X \rightarrow X$ in $\mathcal{C}$ such that $R_I = id_I$, then $Reg_L^1(\mathcal{C})$ is an abelian group under composition.*

(ii) *The set of laycles on $\mathcal{C}$ is a group, denoted by $Z_L^2(\mathcal{C})$.*

(iii) *The map $D^1 : Reg_L^1(\mathcal{C}) \rightarrow Z_L^2(\mathcal{C})$ is a group morphism with image (denoted by $B_L^2(\mathcal{C})$) contained in the centre of $Z_L^2(\mathcal{C})$.*

We denote by $H_L^2(\mathcal{C})$ the group $Z_L^2(\mathcal{C})/B_L^2(\mathcal{C})$, and call it the lazy cohomology of $\mathcal{C}$.

A basic property of lazy cocycles on Hopf algebras (see [1]) is that they act on coquasitriangular structures. This property extends to the categorical setting:

Proposition 2.8 *Let $\mathcal{C}$ be a monoidal category, T a laxycle and c a braiding on $\mathcal{C}$. Then the family $c_{X,Y}^T := T_{Y,X} \circ c_{X,Y} \circ T_{X,Y}^{-1}$ is also a braiding on $\mathcal{C}$.*

Proof. The naturality of c with respect to the morphisms id_X and $T_{Y,Z}^{-1}$ together with (1.6) imply

$$\begin{aligned} (T_{Y,Z}^{-1} \otimes id_X) \circ (id_Y \otimes c_{X,Z}) \circ (c_{X,Y} \otimes id_Z) \\ = (id_Y \otimes c_{X,Z}) \circ (c_{X,Y} \otimes id_Z) \circ (id_X \otimes T_{Y,Z}^{-1}). \end{aligned} \quad (2.10)$$

The naturality of c with respect to the morphisms $T_{X,Y}^{-1}$ and id_Z together with (1.7) imply

$$\begin{aligned} (id_Z \otimes T_{X,Y}^{-1}) \circ (c_{X,Z} \otimes id_Y) \circ (id_X \otimes c_{Y,Z}) \\ = (c_{X,Z} \otimes id_Y) \circ (id_X \otimes c_{Y,Z}) \circ (T_{X,Y}^{-1} \otimes id_Z). \end{aligned} \quad (2.11)$$

We check (1.6) for c^T ; we compute:

$$\begin{aligned} c_{X,Y \otimes Z}^T &= T_{Y \otimes Z, X} \circ c_{X, Y \otimes Z} \circ T_{X, Y \otimes Z}^{-1} \\ &\stackrel{(1.6), (2.2)}{=} (id_Y \otimes T_{Z,X}) \circ T_{Y, Z \otimes X} \circ (T_{Y,Z}^{-1} \otimes id_X) \circ (id_Y \otimes c_{X,Z}) \\ &\quad \circ (c_{X,Y} \otimes id_Z) \circ (id_X \otimes T_{Y,Z}) \circ T_{X \otimes Y, Z}^{-1} \circ (T_{X,Y}^{-1} \otimes id_Z) \\ &\stackrel{(2.10)}{=} (id_Y \otimes T_{Z,X}) \circ T_{Y, Z \otimes X} \circ (id_Y \otimes c_{X,Z}) \\ &\quad \circ (c_{X,Y} \otimes id_Z) \circ T_{X \otimes Y, Z}^{-1} \circ (T_{X,Y}^{-1} \otimes id_Z) \\ &\stackrel{\text{naturality of } T}{=} (id_Y \otimes T_{Z,X}) \circ (id_Y \otimes c_{X,Z}) \circ T_{Y, X \otimes Z} \\ &\quad \circ T_{Y \otimes X, Z}^{-1} \circ (c_{X,Y} \otimes id_Z) \circ (T_{X,Y}^{-1} \otimes id_Z) \\ &\stackrel{(2.2)}{=} (id_Y \otimes T_{Z,X}) \circ (id_Y \otimes c_{X,Z}) \circ (id_Y \otimes T_{X,Z}^{-1}) \\ &\quad \circ (T_{Y,X} \otimes id_Z) \circ (c_{X,Y} \otimes id_Z) \circ (T_{X,Y}^{-1} \otimes id_Z) \\ &= (id_Y \otimes c_{X,Z}^T) \circ (c_{X,Y}^T \otimes id_Z), \quad q.e.d. \end{aligned}$$

Similarly, we check (1.7) for c^T :

$$\begin{aligned} c_{X \otimes Y, Z}^T &= T_{Z, X \otimes Y} \circ c_{X \otimes Y, Z} \circ T_{X \otimes Y, Z}^{-1} \\ &\stackrel{(1.7), (2.2)}{=} (T_{Z,X} \otimes id_Y) \circ T_{Z \otimes X, Y} \circ (id_Z \otimes T_{X,Y}^{-1}) \circ (c_{X,Z} \otimes id_Y) \\ &\quad \circ (id_X \otimes c_{Y,Z}) \circ (T_{X,Y} \otimes id_Z) \circ T_{X, Y \otimes Z}^{-1} \circ (id_X \otimes T_{Y,Z}^{-1}) \\ &\stackrel{(2.11)}{=} (T_{Z,X} \otimes id_Y) \circ T_{Z \otimes X, Y} \circ (c_{X,Z} \otimes id_Y) \\ &\quad \circ (id_X \otimes c_{Y,Z}) \circ T_{X, Y \otimes Z}^{-1} \circ (id_X \otimes T_{Y,Z}^{-1}) \\ &\stackrel{\text{naturality of } T}{=} (T_{Z,X} \otimes id_Y) \circ (c_{X,Z} \otimes id_Y) \circ T_{X \otimes Z, Y} \\ &\quad \circ T_{X, Z \otimes Y}^{-1} \circ (id_X \otimes c_{Y,Z}) \circ (id_X \otimes T_{Y,Z}^{-1}) \\ &\stackrel{(2.2)}{=} (T_{Z,X} \otimes id_Y) \circ (c_{X,Z} \otimes id_Y) \circ (T_{X,Z}^{-1} \otimes id_Y) \\ &\quad \circ (id_X \otimes T_{Z,Y}) \circ (id_X \otimes c_{Y,Z}) \circ (id_X \otimes T_{Y,Z}^{-1}) \\ &= (c_{X,Z}^T \otimes id_Y) \circ (id_X \otimes c_{Y,Z}^T), \end{aligned}$$

finishing the proof. $\square$

Proposition 2.9 *Let $\mathcal{C}$ be a monoidal category, c a braiding on $\mathcal{C}$ and $R_X : X \rightarrow X$ a family of natural isomorphisms in $\mathcal{C}$ such that $R_I = id_I$. Then $c^{D^1(R)} = c$, where $D^1(R)$ is the laycle given by (2.9).*

Proof. Follows immediately by using the naturality of c and R . $\square$

Corollary 2.10 *If $\mathcal{C}$ is a small monoidal category, then the group $H_L^2(\mathcal{C})$ acts on the set of braidings of $\mathcal{C}$.*

Proposition 2.11 *In the hypotheses of Proposition 2.8, the braided monoidal categories $(\mathcal{C}, c)$ and $(\mathcal{C}, c^T)$ are equivalent (as braided monoidal categories).*

Proof. We define the monoidal functor $(F, \varphi_0, \varphi_2) : \mathcal{C} \rightarrow \mathcal{C}$ by $F = id_{\mathcal{C}}$, $\varphi_0 = id_I$ and $\varphi_2(X, Y) : X \otimes Y \rightarrow X \otimes Y$, $\varphi_2(X, Y) := T_{X,Y}^{-1}$, which is obviously a monoidal equivalence. Moreover, the formula $c_{X,Y}^T = T_{Y,X} \circ c_{X,Y} \circ T_{X,Y}^{-1}$ expresses exactly the fact that $(F, \varphi_0, \varphi_2)$ is a braided functor from $(\mathcal{C}, c)$ to $(\mathcal{C}, c^T)$. $\square$

If $\mathcal{C}$ is a braided monoidal category with braiding c , we denote by $Br(\mathcal{C}, c)$ its Brauer group as introduced in [35]. Thus, as a consequence of Proposition 2.11, we obtain the following generalization of [6], Proposition 3.1:

Corollary 2.12 *In the hypotheses of Proposition 2.8, the Brauer groups $Br(\mathcal{C}, c)$ and $Br(\mathcal{C}, c^T)$ are isomorphic.*

Definition 2.13 *Let $\mathcal{C}$ be a monoidal category. A **quasi-braiding** on $\mathcal{C}$ is a family of natural isomorphisms $q_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ in $\mathcal{C}$ satisfying the following axioms (for all $X, Y, Z \in \mathcal{C}$):*

$$q_{I,I} = id_I, \quad (2.12)$$

$$q_{X,Z \otimes Y} \circ (id_X \otimes q_{Y,Z}) = q_{Y \otimes X, Z} \circ (q_{X,Y} \otimes id_Z). \quad (2.13)$$

*If $q_{Y,X} \circ q_{X,Y} = id_{X \otimes Y}$ for all $X, Y \in \mathcal{C}$, then $(\mathcal{C}, q)$ is what Drinfeld calls a **coboundary category** in [13].*

Remark 2.14 *If q is a quasi-braiding on $\mathcal{C}$ then we also have $q_{X,I} = q_{I,X} = id_X$ and*

$$\begin{aligned} (q_{Y,Z} \otimes id_X) \circ q_{X,Y \otimes Z} &= q_{X,Z \otimes Y} \circ (id_X \otimes q_{Y,Z}) \\ &= q_{Y \otimes X, Z} \circ (q_{X,Y} \otimes id_Z) = (id_Z \otimes q_{X,Y}) \circ q_{X \otimes Y, Z}. \end{aligned} \quad (2.14)$$

Consequently, the family $p_{X,Y} := q_{Y,X}^{-1}$ is also a quasi-braiding.

The concept of quasi-braiding was considered (with a different name) by L. M. Ionescu in [18], as follows. Define a monoidal category $\mathcal{C}_{op}$, which is the same as $\mathcal{C}$ as a category, has the same unit I , and reversed tensor product: $X \otimes_{op} Y = Y \otimes X$. Then, a family $q_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ is a quasi-braiding on $\mathcal{C}$ if and only if $(id_{\mathcal{C}}, id_I, \varphi_2(X, Y) := q_{X,Y})$ is a monoidal functor from $\mathcal{C}_{op}$ to $\mathcal{C}$, or equivalently $(id_{\mathcal{C}}, id_I, \varphi_2(X, Y) := q_{Y,X})$ is a monoidal functor from $\mathcal{C}$ to $\mathcal{C}_{op}$. As noted in [18], any braiding is a quasi-braiding (this follows easily by (1.6), (1.7) and (1.8)), and quasi-braidings are related to Drinfeld's coboundary Hopf algebras:

Definition 2.15 ([12]) A coboundary Hopf algebra is a pair (H, R) , where H is a Hopf algebra and $R \in H \otimes H$ is an invertible element such that:

$$R_{12}(\Delta \otimes id)(R) = R_{23}(id \otimes \Delta)(R), \quad (2.15)$$

$$(\varepsilon \otimes id)(R) = (id \otimes \varepsilon)(R) = 1, \quad (2.16)$$

$$\Delta^{cop}(h)R = R\Delta(h), \quad \forall h \in H, \quad (2.17)$$

$$R_{21}R = 1 \otimes 1. \quad (2.18)$$

If R does not necessarily satisfy (2.18), we call it a **quasi-coboundary**.

Proposition 2.16 ([18]) Let H be a Hopf algebra and $R = R^1 \otimes R^2 \in H \otimes H$ an invertible element. If U, V are left H -modules, define $q_{U,V} : U \otimes V \rightarrow V \otimes U$ by $q_{U,V}(u \otimes v) = R^2 \cdot v \otimes R^1 \cdot u$. Then q is a quasi-braiding on ${}_H\mathcal{M}$ if and only if R is a quasi-coboundary on H .

Remark 2.17 If T is a laycle on a monoidal category $\mathcal{C}$, then the family $\mathcal{T}_{X,Y} := T_{Y,X}$ is a laycle on $\mathcal{C}_{op}$.

From the description of laycles and quasi-braidings as monoidal structures for some identity functors and the fact that a composition of monoidal functors is monoidal, we obtain:

Proposition 2.18 Let $\mathcal{C}$ be a monoidal category, T a laycle and p, q two quasi-braidings on $\mathcal{C}$. Then the family $D_{X,Y} := p_{Y,X} \circ q_{X,Y}$ is a laycle on $\mathcal{C}$ and the families $q'_{X,Y} := T_{Y,X} \circ q_{X,Y}$ and $q''_{X,Y} := q_{X,Y} \circ T_{X,Y}$ are quasi-braidings on $\mathcal{C}$.

Corollary 2.19 Let $\mathcal{C}$ be a monoidal category, T a laycle and q a quasi-braiding on $\mathcal{C}$. Then the family $q_{X,Y}^T := T_{Y,X} \circ q_{X,Y} \circ T_{X,Y}^{-1}$ is also a quasi-braiding on $\mathcal{C}$.

Remark 2.20 Proposition 2.9 is also true with quasi-braidings instead of braidings, so we obtain also an action of $H_L^2(\mathcal{C})$ on the set of quasi-braidings of $\mathcal{C}$.

Proposition 2.21 Let $\mathcal{C}$ be a monoidal category, c a braiding and q a quasi-braiding on $\mathcal{C}$. Then the family $c_{X,Y}^q := q_{Y,X}^{-1} \circ c_{Y,X} \circ q_{X,Y}$ is also a braiding on $\mathcal{C}$. Moreover, the braided categories $(\mathcal{C}, c)$ and $(\mathcal{C}, c^q)$ are braided equivalent.

Proof. Follows immediately from Proposition 2.8, since $c^q = c^T$, where T is the laycle $T_{X,Y} = q_{X,Y}^{-1} \circ c_{X,Y}$. $\square$

Remark 2.22 For the particular case when q itself is a braiding, we will obtain an alternative proof in Proposition 5.3.

Let now $\mathcal{C}$ be a small monoidal category. We denote by $\mathbb{Z}^2(\mathcal{C})$ the set of all natural isomorphisms in $\mathcal{C}$ that are laycles or quasi-braidings. Then, with notation as in Proposition 2.7, we have:

Proposition 2.23 (i) $\mathbb{Z}^2(\mathcal{C})$ is a group.
(ii) $Z_L^2(\mathcal{C})$ is an index 2 subgroup in $\mathbb{Z}^2(\mathcal{C})$.
(iii) $B_L^2(\mathcal{C})$ is a central subgroup in $\mathbb{Z}^2(\mathcal{C})$.

We define the “cohomology group” $\mathbb{H}^2(\mathcal{C}) := \mathbb{Z}^2(\mathcal{C})/B_L^2(\mathcal{C})$.

Proof. We give first the explicit description of the multiplication in $\mathbb{Z}^2(\mathcal{C})$. Take R and P quasi-braidings, S and T laycles. We have, for all $U, V \in \mathcal{C}$:

$$\begin{aligned}(ST)_{U,V} &= S_{U,V} \circ T_{U,V}, \\ (TR)_{U,V} &= T_{V,U} \circ R_{U,V}, \\ (RT)_{U,V} &= R_{U,V} \circ T_{U,V}, \\ (RP)_{U,V} &= R_{V,U} \circ P_{U,V}.\end{aligned}$$

Now (i) and (ii) follow from Proposition 2.18, while (iii) is just an easy computation. $\square$

Similarly, if H is a Hopf algebra, we may consider the group $\mathbb{Z}^2(H)$ consisting of the elements in $H \otimes H$ that are lazy twists or quasi-coboundaries, its central subgroup $B_{LT}^2(H)$ and the “cohomology group” $\mathbb{H}^2(H) = \mathbb{Z}^2(H)/B_{LT}^2(H)$.

Example 2.24 Let k be a field with $\text{char}(k) \neq 2$ and $H = k[C_2]$, the group algebra of the cyclic group with two elements C_2 (denote its generator by g). One can see that the lazy twists on H are given by the formula $T_a = \frac{3+a}{4}(1 \otimes 1) + \frac{1-a}{4}(1 \otimes g) + \frac{1-a}{4}(g \otimes 1) - \frac{1-a}{4}(g \otimes g)$, with $a \in k^*$. It is interesting to note that T_0 is not invertible but has all the other properties in the definition of a lazy twist.

Consider the element $\theta_\alpha = \frac{1+g}{2} + \alpha \frac{1-g}{2} \in H$, with $\alpha \in k$. One can see that θ_α is invertible if and only if $\alpha \neq 0$. Also it is easy to see that $T_{\alpha^{-2}} = \Delta(\theta_\alpha)(\theta_\alpha^{-1} \otimes \theta_\alpha^{-1})$ and so T_a is trivial in $\mathbb{H}^2(H)$ if and only if $a \in (k^*)^2$. One can also note that $T_a T_b = T_{ab}$.

Since H is commutative and cocommutative, one can see that the quasi-coboundaries for H are given by the formula $R_a = \frac{3+a}{4}(1 \otimes 1) + \frac{1-a}{4}(1 \otimes g) + \frac{1-a}{4}(g \otimes 1) - \frac{1-a}{4}(g \otimes g)$, with $a \in k^*$. Among these, only R_1 and R_{-1} are quasitriangular. If we put everything together we obtain $\mathbb{H}^2(H) = k^*/(k^*)^2 \times C_2$.

3 Strong twines and pseudosymmetric braidings

A key result for this section is the following characterization of strong twines:

Proposition 3.1 *Let $\mathcal{C}$ be a monoidal category and T a laycle on $\mathcal{C}$. Then T is a strong twine if and only if the families T^b and T^f given by (2.3) and (2.4) coincide.*

Proof. Let $X, Y, Z \in \mathcal{C}$ and assume that T is a strong twine; then we have:

$$\begin{aligned}T_{X,Y,Z}^b &= (id_X \otimes T_{Y,Z}^{-1}) \circ T_{X \otimes Y, Z} \\ &= (T_{X,Y}^{-1} \otimes id_Z) \circ (T_{X,Y} \otimes id_Z) \circ (id_X \otimes T_{Y,Z}^{-1}) \circ T_{X \otimes Y, Z} \\ &\stackrel{(1.20)}{=} (T_{X,Y}^{-1} \otimes id_Z) \circ (id_X \otimes T_{Y,Z}^{-1}) \circ (T_{X,Y} \otimes id_Z) \circ T_{X \otimes Y, Z} \\ &\stackrel{(1.19)}{=} (T_{X,Y}^{-1} \otimes id_Z) \circ (id_X \otimes T_{Y,Z}^{-1}) \circ (id_X \otimes T_{Y,Z}) \circ T_{X,Y \otimes Z} \\ &= (T_{X,Y}^{-1} \otimes id_Z) \circ T_{X,Y \otimes Z} \\ &= T_{X,Y,Z}^f.\end{aligned}$$

Conversely, assume that $T^b = T^f$. By using (2.3), (2.2) and (2.4) it is easy to see that $T_{X,Y,Z}^b \circ (T_{X,Y} \otimes id_Z) \circ (id_X \otimes T_{Y,Z}) = T_{X,Y,Z}^f \circ (id_X \otimes T_{Y,Z}) \circ (T_{X,Y} \otimes id_Z)$, and since $T^b = T^f$ it follows that (1.20) holds. $\square$

Definition 3.2 ([2]) Let $\mathcal{C}$ be a monoidal category. A D -structure on $\mathcal{C}$ consists of a family of natural morphisms $R_X : X \rightarrow X$ in $\mathcal{C}$, such that $R_I = id_I$ and (for all $X, Y, Z \in \mathcal{C}$):

$$(R_{X \otimes Y} \otimes id_Z)(id_X \otimes R_{Y \otimes Z}) = (id_X \otimes R_{Y \otimes Z})(R_{X \otimes Y} \otimes id_Z). \quad (3.1)$$

It was proved in [29] that if R is a D -structure consisting of isomorphisms then the family $D^1(R)$ given by (2.9) is a strong twine. Using Proposition 3.1 we can prove the converse:

Proposition 3.3 Let $\mathcal{C}$ be a monoidal category and $R_X : X \rightarrow X$ a family of natural isomorphisms in $\mathcal{C}$ with $R_I = id_I$. Then $D^1(R)$ is a strong twine if and only if R is a D -structure.

Proof. We compute:

$$\begin{aligned} D^1(R)_{X,Y,Z}^b &= (id_X \otimes R_{Y \otimes Z}) \circ (id_X \otimes R_Y^{-1} \otimes R_Z^{-1}) \circ (R_{X \otimes Y} \otimes R_Z) \circ R_{X \otimes Y \otimes Z}^{-1} \\ &= (id_X \otimes R_{Y \otimes Z}) \circ (id_X \otimes R_Y^{-1} \otimes R_Z^{-1}) \circ (id_X \otimes R_Y \otimes R_Z) \\ &\quad \circ (id_X \otimes R_Y^{-1} \otimes id_Z) \circ (R_{X \otimes Y} \otimes id_Z) \circ R_{X \otimes Y \otimes Z}^{-1} \\ &= (id_X \otimes R_{Y \otimes Z}) \circ (id_X \otimes R_Y^{-1} \otimes id_Z) \circ (R_{X \otimes Y} \otimes id_Z) \circ R_{X \otimes Y \otimes Z}^{-1} \\ &\stackrel{\text{naturality of } R}{=} (id_X \otimes R_Y^{-1} \otimes id_Z) \circ (id_X \otimes R_{Y \otimes Z}) \circ (R_{X \otimes Y} \otimes id_Z) \circ R_{X \otimes Y \otimes Z}^{-1}, \end{aligned}$$

and similarly one can see that

$$D^1(R)_{X,Y,Z}^f = (id_X \otimes R_Y^{-1} \otimes id_Z) \circ (R_{X \otimes Y} \otimes id_Z) \circ (id_X \otimes R_{Y \otimes Z}) \circ R_{X \otimes Y \otimes Z}^{-1},$$

and it is clear that $D^1(R)^b = D^1(R)^f$ (i.e. $D^1(R)$ is a strong twine) if and only if (3.1) holds. $\square$

We recall that a (generalized) double braiding is always a twine; it is natural to ask under what conditions is it a strong twine. The answer is provided by our next result:

Theorem 3.4 Let $\mathcal{C}$ be a monoidal category, c and d braidings on $\mathcal{C}$ and $T_{X,Y} = d_{Y,X} \circ c_{X,Y}$. Then T is a strong twine if and only if the following relation holds, for all $X, Y, Z \in \mathcal{C}$:

$$\begin{aligned} (d_{Z,X} \otimes id_Y) \circ (id_Z \otimes c_{X,Y}^{-1}) \circ (c_{Y,Z} \otimes id_X) \\ \circ (d_{Z,Y} \otimes id_X) \circ (id_Z \otimes c_{X,Y}) \circ (c_{X,Z} \otimes id_Y) \\ = (id_X \otimes c_{Y,Z}) \circ (d_{X,Y}^{-1} \otimes id_Z) \circ (id_Y \otimes d_{Z,X}) \\ \circ (id_Y \otimes c_{X,Z}) \circ (d_{X,Y} \otimes id_Z) \circ (id_X \otimes d_{Z,Y}). \quad (3.2) \end{aligned}$$

Proof. We compute the families T^b and T^f :

$$\begin{aligned} T_{X,Y,Z}^b &= (id_X \otimes T_{Y,Z}^{-1}) \circ T_{X \otimes Y, Z} \\ &\stackrel{(1.6)}{=} (id_X \otimes c_{Y,Z}^{-1}) \circ (id_X \otimes d_{Z,Y}^{-1}) \circ (id_X \otimes d_{Z,Y}) \circ (d_{Z,X} \otimes id_Y) \circ c_{X \otimes Y, Z} \\ &= (id_X \otimes c_{Y,Z}^{-1}) \circ (d_{Z,X} \otimes id_Y) \circ c_{X \otimes Y, Z}, \\ T_{X,Y,Z}^f &= T_{X \otimes Y, Z} \circ (id_X \otimes T_{Y,Z}^{-1}) \\ &\stackrel{(1.7)}{=} d_{Z,X \otimes Y} \circ (c_{X,Z} \otimes id_Y) \circ (id_X \otimes c_{Y,Z}) \circ (id_X \otimes c_{Y,Z}^{-1}) \circ (id_X \otimes d_{Z,Y}^{-1}) \\ &= d_{Z,X \otimes Y} \circ (c_{X,Z} \otimes id_Y) \circ (id_X \otimes d_{Z,Y}^{-1}). \end{aligned}$$

By Proposition 3.1, T is a strong twine if and only if $T^b = T^f$, and this holds if and only if

$$(d_{Z,X} \otimes id_Y) \circ c_{X \otimes Y, Z} \circ (id_X \otimes d_{Z,Y}) = (id_X \otimes c_{Y,Z}) \circ d_{Z, X \otimes Y} \circ (c_{X,Z} \otimes id_Y). \quad (3.3)$$

Thus, it is enough to prove that the left hand sides of equations (3.2) and (3.3) coincide, and the same for the right hand sides. We compute:

$$\begin{aligned} & (d_{Z,X} \otimes id_Y) \circ c_{X \otimes Y, Z} \circ (id_X \otimes d_{Z,Y}) \\ &= (d_{Z,X} \otimes id_Y) \circ (id_Z \otimes c_{X,Y}^{-1}) \circ (c_{Y,Z} \otimes id_X) \circ c_{X, Y \otimes Z} \circ (id_X \otimes d_{Z,Y}) \\ &= (d_{Z,X} \otimes id_Y) \circ (id_Z \otimes c_{X,Y}^{-1}) \circ (c_{Y,Z} \otimes id_X) \circ (d_{Z,Y} \otimes id_X) \circ c_{X, Z \otimes Y} \\ &= (d_{Z,X} \otimes id_Y) \circ (id_Z \otimes c_{X,Y}^{-1}) \circ (c_{Y,Z} \otimes id_X) \\ &\quad \circ (d_{Z,Y} \otimes id_X) \circ (id_Z \otimes c_{X,Y}) \circ (c_{X,Z} \otimes id_Y) \end{aligned}$$

(for the first equality we used (2.14), for the second the naturality of c and for the third (1.6)),

$$\begin{aligned} & (id_X \otimes c_{Y,Z}) \circ d_{Z, X \otimes Y} \circ (c_{X,Z} \otimes id_Y) \\ &= (id_X \otimes c_{Y,Z}) \circ (d_{X,Y}^{-1} \otimes id_Z) \circ (id_Y \otimes d_{Z,X}) \circ d_{Z \otimes X, Y} \circ (c_{X,Z} \otimes id_Y) \\ &= (id_X \otimes c_{Y,Z}) \circ (d_{X,Y}^{-1} \otimes id_Z) \circ (id_Y \otimes d_{Z,X}) \circ (id_Y \otimes c_{X,Z}) \circ d_{X \otimes Z, Y} \\ &= (id_X \otimes c_{Y,Z}) \circ (d_{X,Y}^{-1} \otimes id_Z) \circ (id_Y \otimes d_{Z,X}) \\ &\quad \circ (id_Y \otimes c_{X,Z}) \circ (d_{X,Y} \otimes id_Z) \circ (id_X \otimes d_{Z,Y}) \end{aligned}$$

(for the first equality we used (2.14), for the second the naturality of d and for the third (1.7)), finishing the proof. $\square$

Definition 3.5 Let $\mathcal{C}$ be a monoidal category and c a braiding on $\mathcal{C}$. We will say that c is a **pseudosymmetry** if the following condition holds, for all $X, Y, Z \in \mathcal{C}$:

$$\begin{aligned} & (c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1}) \circ (c_{X,Y} \otimes id_Z) \\ &= (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y) \circ (id_X \otimes c_{Y,Z}). \quad (3.4) \end{aligned}$$

In this case we will say that $\mathcal{C}$ is a **pseudosymmetric braided category**.

If c is a symmetry, i.e. $c_{Z,X}^{-1} = c_{X,Z}$, then obviously c is a pseudosymmetry, by (1.8).

Theorem 3.6 Let $\mathcal{C}$ be a monoidal category and c a braiding on $\mathcal{C}$. Then the double braiding $T_{X,Y} = c_{Y,X} \circ c_{X,Y}$ is a strong twine if and only if c is a pseudosymmetry.

Proof. In (3.2) written for $c = d$ we have, by (1.8),

$$(c_{Z,Y} \otimes id_X) \circ (id_Z \otimes c_{X,Y}) \circ (c_{X,Z} \otimes id_Y) = (id_Y \otimes c_{X,Z}) \circ (c_{X,Y} \otimes id_Z) \circ (id_X \otimes c_{Z,Y}),$$

so (3.2) reduces in this case to (3.4). $\square$

Let H be a Hopf algebra. Consider the category ${}_H\mathcal{YD}^H$ of left-right Yetter-Drinfeld modules over H , whose objects are vector spaces M that are left H -modules (denote the action by $h \otimes m \mapsto h \cdot m$) and right H -comodules (denote the coaction by $m \mapsto m_{(0)} \otimes m_{(1)} \in M \otimes H$) satisfying the compatibility condition

$$(h \cdot m)_{(0)} \otimes (h \cdot m)_{(1)} = h_2 \cdot m_{(0)} \otimes h_3 m_{(1)} S^{-1}(h_1), \quad \forall h \in H, m \in M. \quad (3.5)$$

It is a monoidal category, with tensor product given by

$$h \cdot (m \otimes n) = h_1 \cdot m \otimes h_2 \cdot n, \quad (m \otimes n)_{(0)} \otimes (m \otimes n)_{(1)} = m_{(0)} \otimes n_{(0)} \otimes n_{(1)} m_{(1)}.$$

Moreover, it has a (canonical) braiding given by

$$\begin{aligned} c_{M,N} : M \otimes N &\rightarrow N \otimes M, & c_{M,N}(m \otimes n) &= n_{(0)} \otimes n_{(1)} \cdot m, \\ c_{M,N}^{-1} : N \otimes M &\rightarrow M \otimes N, & c_{M,N}^{-1}(n \otimes m) &= S(n_{(1)}) \cdot m \otimes n_{(0)}. \end{aligned}$$

It is known (cf. [30]) that this braiding is a symmetry only in the degenerate case $H = k$.

Theorem 3.7 *The canonical braiding of ${}_H\mathcal{YD}^H$ is pseudosymmetric if and only if H is commutative and cocommutative.*

Proof. Assume first that H is commutative and cocommutative; in this case, the compatibility condition (3.5) becomes the Long condition

$$(h \cdot m)_{(0)} \otimes (h \cdot m)_{(1)} = h \cdot m_{(0)} \otimes m_{(1)}, \quad \forall h \in H, m \in M. \quad (3.6)$$

For all $X, Y, Z \in {}_H\mathcal{YD}^H$, $x \in X$, $y \in Y$, $z \in Z$ we compute:

$$\begin{aligned} & (c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1}) \circ (c_{X,Y} \otimes id_Z)(x \otimes y \otimes z) \\ &= (c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1})(y_{(0)} \otimes y_{(1)} \cdot x \otimes z) \\ &= (c_{Y,Z} \otimes id_X)(y_{(0)} \otimes S((y_{(1)} \cdot x)_{(1)}) \cdot z \otimes (y_{(1)} \cdot x)_{(0)}) \\ &\stackrel{(3.6)}{=} (c_{Y,Z} \otimes id_X)(y_{(0)} \otimes S(x_{(1)}) \cdot z \otimes y_{(1)} \cdot x_{(0)}) \\ &= (S(x_{(1)}) \cdot z)_{(0)} \otimes (S(x_{(1)}) \cdot z)_{(1)} \cdot y_{(0)} \otimes y_{(1)} \cdot x_{(0)} \\ &\stackrel{(3.6)}{=} S(x_{(1)}) \cdot z_{(0)} \otimes z_{(1)} \cdot y_{(0)} \otimes y_{(1)} \cdot x_{(0)} \\ &\stackrel{(3.6)}{=} S(x_{(1)}) \cdot z_{(0)} \otimes (z_{(1)} \cdot y)_{(0)} \otimes (z_{(1)} \cdot y)_{(1)} \cdot x_{(0)} \\ &= (id_Z \otimes c_{X,Y})(S(x_{(1)}) \cdot z_{(0)} \otimes x_{(0)} \otimes z_{(1)} \cdot y) \\ &= (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y)(x \otimes z_{(0)} \otimes z_{(1)} \cdot y) \\ &= (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y) \circ (id_X \otimes c_{Y,Z})(x \otimes y \otimes z), \end{aligned}$$

proving that c is pseudosymmetric.

Conversely, assume that c is pseudosymmetric. We consider the two usual Yetter-Drinfeld structures on the vector space H : the first one, denoted by H_1 , is H with the usual (regular) left module structure and with comodule structure $\rho_1(h) = h_2 \otimes h_3 S^{-1}(h_1)$, and the second, denoted by H_2 , is H with module structure given by $h \cdot g = h_2 g S^{-1}(h_1)$ and comodule structure $\rho_2(h) = h_1 \otimes h_2$.

We prove first that H is cocommutative. Let $h \in H$; we will apply the pseudosymmetry condition (3.4) for $X = H_1$, $Y = H_2$, $Z = H_1$ on the element $1 \otimes h \otimes 1$:

$$\begin{aligned} & (c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1}) \circ (c_{X,Y} \otimes id_Z)(1 \otimes h \otimes 1) \\ &= (c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1})(h_1 \otimes h_2 \otimes 1) \\ &= (c_{Y,Z} \otimes id_X)(h_1 \otimes h_2 S(h_4) \otimes h_3) \\ &= (h_2 S(h_4))_2 \otimes [(h_2 S(h_4))_3 S^{-1}((h_2 S(h_4))_1)] \cdot h_1 \otimes h_3, \end{aligned}$$

$$\begin{aligned}
& (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y) \circ (id_X \otimes c_{Y,Z})(1 \otimes h \otimes 1) \\
&= (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y)(1 \otimes 1 \otimes h) \\
&= (id_Z \otimes c_{X,Y})(1 \otimes 1 \otimes h) \\
&= 1 \otimes h_1 \otimes h_2,
\end{aligned}$$

so we obtain

$$(h_2 S(h_4))_2 \otimes [(h_2 S(h_4))_3 S^{-1}((h_2 S(h_4))_1)] \cdot h_1 \otimes h_3 = 1 \otimes h_1 \otimes h_2.$$

By applying $id \otimes \varepsilon \otimes id$ we get $h_1 S(h_3) \otimes h_2 = 1 \otimes h$, which, by making convolution with $S(h) \otimes 1$, becomes $S(h_1) h_2 S(h_4) \otimes h_3 = S(h_1) \otimes h_2$, and so we obtain $S(h_2) \otimes h_1 = S(h_1) \otimes h_2$, which implies $\Delta^{cop}(h) = \Delta(h)$, i.e. H is cocommutative.

We prove now that H is commutative. Note first that cocommutativity implies $c_{H_2, H_1}(b \otimes a) = a \otimes b$, for all $a, b \in H$. Let now $g, h \in H$; we will apply the pseudosymmetry condition (3.4) for $X = H_1$, $Y = H_2$, $Z = H_2$ on the element $1 \otimes g \otimes h$:

$$\begin{aligned}
& (c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1}) \circ (c_{X,Y} \otimes id_Z)(1 \otimes g \otimes h) \\
&= (c_{Y,Z} \otimes id_X) \circ (id_Y \otimes c_{Z,X}^{-1})(g_1 \otimes g_2 \otimes h) \\
&= (c_{Y,Z} \otimes id_X)(g_1 \otimes h \otimes g_2) \\
&= h_1 \otimes h_3 g_1 S^{-1}(h_2) \otimes g_2, \\
& (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y) \circ (id_X \otimes c_{Y,Z})(1 \otimes g \otimes h) \\
&= (id_Z \otimes c_{X,Y}) \circ (c_{Z,X}^{-1} \otimes id_Y)(1 \otimes h_1 \otimes h_3 g S^{-1}(h_2)) \\
&= (id_Z \otimes c_{X,Y})(h_1 \otimes 1 \otimes h_3 g S^{-1}(h_2)) \\
&= h_1 \otimes (h_3 g S^{-1}(h_2))_1 \otimes (h_3 g S^{-1}(h_2))_2,
\end{aligned}$$

and so we obtain

$$h_1 \otimes h_3 g_1 S^{-1}(h_2) \otimes g_2 = h_1 \otimes (h_3 g S^{-1}(h_2))_1 \otimes (h_3 g S^{-1}(h_2))_2.$$

By applying $id \otimes \varepsilon \otimes id$ we get $h_1 \otimes h_3 g S^{-1}(h_2) = h \otimes g$, which implies $h_3 g S^{-1}(h_2) h_1 = gh$, that is $hg = gh$ and hence H is commutative. $\square$

Corollary 3.8 *For H a commutative and cocommutative Hopf algebra, the double braiding*

$$T_{X,Y}(x \otimes y) = (c_{Y,X} \circ c_{X,Y})(x \otimes y) = y_{(1)} \cdot x_{(0)} \otimes x_{(1)} \cdot y_{(0)}$$

is a strong twine on ${}_H \mathcal{YD}^H$.

Definition 3.9 *If H is a Hopf algebra and $R \in H \otimes H$ is a quasitriangular structure, we will say that R is **pseudotriangular** if*

$$R_{12} R_{31}^{-1} R_{23} = R_{23} R_{31}^{-1} R_{12}. \quad (3.7)$$

If R is a pseudotriangular structure then it is easy to see that the braiding on ${}_H \mathcal{M}$ given by $c_{M,N} : M \otimes N \rightarrow N \otimes M$, $c_{M,N}(m \otimes n) = R^2 \cdot n \otimes R^1 \cdot m$, is pseudosymmetric. Also, it is obvious that if R is triangular (i.e. $R_{21} R = 1 \otimes 1$) then R is pseudotriangular, because in this case (3.7) becomes the quantum Yang-Baxter equation $R_{12} R_{13} R_{23} = R_{23} R_{13} R_{12}$. We have also the Hopf-algebraic counterpart of Theorem 3.6:

Proposition 3.10 *Let (H, R) be a quasitriangular Hopf algebra. Then R is pseudotriangular if and only if the lazy twist $F = R_{21}R$ satisfies the condition $F_{12}F_{23} = F_{23}F_{12}$ (i.e. F is neat, in the terminology of [29]).*

Example 3.11 If H is a commutative Hopf algebra, then any quasitriangular structure on H is pseudotriangular. For instance, if k has characteristic zero and contains a primitive root of unity of degree n , then the group algebra of the cyclic group $\mathbb{Z}_n$ admits a certain quasitriangular structure (constructed in [25], [31]) which is *not* triangular for $n \geq 3$. Thus, for $n \geq 3$, the category of representations of $\mathbb{Z}_n$ admits a pseudosymmetric braiding which is not symmetric.

Remark 3.12 Let H be a finite dimensional Hopf algebra. It is well-known that the category of Yetter-Drinfeld modules ${}_H\mathcal{YD}^H$ is braided equivalent to the category ${}_{D(H)}\mathcal{M}$ of left modules over the Drinfeld double of H (realized on $H^{*cop} \otimes H$ and with quasitriangular structure given by $R = \sum(\varepsilon \otimes e_i) \otimes (e^i \otimes 1)$, where $\{e_i\}, \{e^i\}$ are dual bases in H and H^*). Thus, via Theorem 3.7, we obtain that R is pseudotriangular if and only if H is commutative and cocommutative. In particular, if G is a finite, noncommutative group then $(D(k[G]), R)$ is quasitriangular but not pseudotriangular.

Definition 3.13 ([23]) *Let (H, R) be a quasitriangular Hopf algebra. The element R is called **almost-triangular** if $R_{21}R$ is central in $H \otimes H$.*

Remark 3.14 *By Proposition 3.10 it follows that an almost-triangular structure is pseudotriangular. The converse is not true, a counterexample is provided by Proposition 3.15 below.*

Assume now that $\text{char}(k) \neq 2$ and consider the 2^{n+1} -dimensional Hopf algebra $E(n)$ generated by $c, x_1, \dots, x_n$ with relations $c^2 = 1, x_i^2 = 0, x_i c + c x_i = 0$ and $x_i x_j + x_j x_i = 0$, for all $i, j \in \{1, \dots, n\}$, and coalgebra structure $\Delta(c) = c \otimes c, \Delta(x_i) = 1 \otimes x_i + x_i \otimes c$, for all $i \in \{1, \dots, n\}$. The quasitriangular structures of $E(n)$ have been classified in [27], they are in bijection with $n \times n$ matrices with entries in k , and moreover the quasitriangular structure R_A corresponding to the matrix A is given by an explicit formula, generalizing the cases $n = 1$ from [32] and $n = 2$ from [17]. By [27] and [8] we know that R_A is triangular if and only if the matrix A is symmetric.

Proposition 3.15 *For any $n \times n$ matrix A , the quasitriangular structure R_A is pseudotriangular, and it is almost-triangular if and only if A is symmetric (thus the only almost-triangular structures of $E(n)$ are the triangular ones).*

Proof. We present first an alternative description for the quasitriangular structure R_A . For every $a \in k$ and $i, j \in \{1, \dots, n\}$ we define the element

$$T_{i,j}(a) := 1 \otimes 1 + a(x_i \otimes c x_j) \in E(n) \otimes E(n). \quad (3.8)$$

It is easy to see that $T_{i,j}(a)$ is a lazy twist, $T_{i,j}(a)T_{i,j}(b) = T_{i,j}(a+b)$ and $T_{i,j}(a)T_{k,l}(b) = T_{k,l}(b)T_{i,j}(a)$, for all $a, b \in k$ and $i, j, k, l \in \{1, \dots, n\}$. If $A = (a_{ij})_{i,j=1, \dots, n}$ is an $n \times n$ matrix, we define the element

$$T_A := \prod_{i,j=1}^n T_{i,j}(a_{ij}) \in E(n) \otimes E(n) \quad (3.9)$$

(note that the order of the factors does not matter since they all commute). It is clear that if B is another $n \times n$ matrix then $T_A T_B = T_{A+B}$. One can also see that the element T_A is given by the formula

$$T_A = 1 \otimes 1 + \sum_{|P|=|F|} (-1)^{\frac{|P|(|P|-1)}{2}} \det(P, F) x_P \otimes c^{|P|} x_F, \quad (3.10)$$

where the sum is made over all nonempty subsets P, F of $\{1, \dots, n\}$ such that $|P| = |F|$, and if $P = \{i_1 < i_2 < \dots < i_s\}$ and $F = \{j_1 < j_2 < \dots < j_s\}$ then $\det(P, F)$ is the determinant of the $s \times s$ matrix obtained at the intersection of the rows $i_1, \dots, i_s$ and columns $j_1, \dots, j_s$ of the matrix A , and $x_P = x_{i_1} \cdots x_{i_s}$, $x_F = x_{j_1} \cdots x_{j_s}$. In particular we obtain $T_0 = 1 \otimes 1$ and $T_A^{-1} = T_{-A}$.

Define now the element

$$R := \frac{1}{2}(1 \otimes 1 + c \otimes 1 + 1 \otimes c - c \otimes c) \in E(n) \otimes E(n),$$

which is a triangular structure for $E(n)$. From the formula for the quasitriangular structure R_A in [27] and (3.10) we immediately obtain

$$R_A = R T_A. \quad (3.11)$$

If we denote by A^t the transpose of a matrix A , then we know from [8] that

$$R_A^{-1} = (R_{A^t})_{21}, \quad (3.12)$$

a consequence of which is the relation $(R_A)_{21} R_B = T_{B-A^t}$, for any $n \times n$ matrices A and B . We record also the obvious relation $R_A T_B = R_{A+B}$, as well as $(T_A)_{21} R_B = R_{B-A^t}$.

Let now A be an $n \times n$ matrix; we will prove that R_A is pseudotriangular. In view of (3.12), what we need to prove is the relation

$$(R_A)_{12} (R_{A^t})_{13} (R_A)_{23} = (R_A)_{23} (R_{A^t})_{13} (R_A)_{12}. \quad (3.13)$$

We will actually prove something more general, namely

$$(R_A)_{12} (R_B)_{13} (R_C)_{23} = (R_C)_{23} (R_B)_{13} (R_A)_{12}, \quad (3.14)$$

for any $n \times n$ matrices A, B and C . We introduce the following notation, for $a \in k$ and $i, j \in \{1, \dots, n\}$:

$$\begin{aligned} T_{i,j}(a)_{12c} &:= 1 \otimes 1 \otimes 1 + a x_i \otimes c x_j \otimes c, \\ T_{i,j}(a)_{1c3} &:= 1 \otimes 1 \otimes 1 + a x_i \otimes c \otimes c x_j, \\ T_{i,j}(a)_{c23} &:= 1 \otimes 1 \otimes 1 + c \otimes a x_i \otimes c x_j. \end{aligned}$$

By direct computation one can prove the following relations:

$$\begin{aligned} T_{i,j}(a)_{23} R_{13} &= R_{13} T_{i,j}(a)_{c23}, \\ T_{i,j}(a)_{c23} R_{12} &= R_{12} T_{i,j}(a)_{23}, \\ T_{i,j}(a)_{13} R_{12} &= R_{12} T_{i,j}(a)_{1c3}, \\ T_{i,j}(a)_{13} R_{23} &= R_{23} T_{i,j}(a)_{1c3}, \\ T_{i,j}(a)_{12} R_{13} &= R_{13} T_{i,j}(a)_{12c}, \\ T_{i,j}(a)_{12c} R_{23} &= R_{23} T_{i,j}(a)_{12}. \end{aligned}$$

One can also see that, for all $i, j, k, l, p, q \in \{1, \dots, n\}$ and $x, y, z \in k$, all the elements $T_{i,j}(x)_{23}$, $T_{k,l}(y)_{12}$ and $T_{p,q}(z)_{1c3}$ commute with each other. Using all these facts together with the formulae (3.11) and (3.9) we obtain

$$\begin{aligned}(R_A)_{12}(R_B)_{13}(R_C)_{23} &= R_{12}R_{13}R_{23}(T_A)_{12}(T_B)_{1c3}(T_C)_{23}, \\ (R_C)_{23}(R_B)_{13}(R_A)_{12} &= R_{23}R_{13}R_{12}(T_C)_{23}(T_B)_{1c3}(T_A)_{12},\end{aligned}$$

and the right hand sides are equal because of the above-mentioned commutation relations together with the fact that R satisfies the Yang-Baxter equation.

We prove now that R_A is almost-triangular if and only if A is symmetric. Let B be an $n \times n$ matrix; it is easy to see that T_B is central in $E(n) \otimes E(n)$ if and only if $B = 0$, because if $B \neq 0$ then T_B does not commute with $1 \otimes c$. We have seen above that $(R_A)_{21}R_A = T_{A-A^t}$, and so $(R_A)_{21}R_A$ is central if and only if $A = A^t$. $\square$

Remark 3.16 We consider the group $\mathbb{Z}^2(E(n))$ as in Section 2, and inside it the set $G_n := \{T_A, R_A\}$, where A is an $n \times n$ matrix. If we denote by $*$ the multiplication in $\mathbb{Z}^2(E(n))$, then we have

$$\begin{aligned}T_A * T_B &= T_A T_B = T_{A+B}, \\ R_A * T_B &= R_A T_B = R_{A+B}, \\ T_A * R_B &= (T_A)_{21} R_B = R_{B-A^t}, \\ R_A * R_B &= (R_A)_{21} R_B = T_{B-A^t},\end{aligned}$$

and so G_n is a subgroup of $\mathbb{Z}^2(E(n))$ (note that the inverse of R_A in this group is R_{A^t}). The above formulae imply $G_n \simeq \mathbb{Z}_2 \ltimes (M_n(k), +)$, a semidirect product, where the action of $\mathbb{Z}_2$ on $(M_n(k), +)$ is given by $A \cdot g = -A^t$ (g is the generator of $\mathbb{Z}_2$), and the correspondence is given by $T_A \mapsto (1, A)$, $R_A \mapsto (g, A)$. For $n = 1$ ($E(1)$ is Sweedler's 4-dimensional Hopf algebra), one can prove by direct computation that $G_1 = \mathbb{Z}^2(E(1))$.

4 Laycles, pseudotwistors and R-matrices

We recall the following concept and result from [24]:

Proposition 4.1 ([24]) *Let $\mathcal{C}$ be a monoidal category, A an algebra in $\mathcal{C}$ with multiplication μ and unit u , $T : A \otimes A \rightarrow A \otimes A$ a morphism in $\mathcal{C}$ such that $T \circ (u \otimes id_A) = u \otimes id_A$ and $T \circ (id_A \otimes u) = id_A \otimes u$. Assume that there exist two morphisms $\tilde{T}_1, \tilde{T}_2 : A \otimes A \otimes A \rightarrow A \otimes A \otimes A$ in $\mathcal{C}$ such that*

$$(id_A \otimes \mu) \circ \tilde{T}_1 \circ (T \otimes id_A) = T \circ (id_A \otimes \mu), \quad (4.1)$$

$$(\mu \otimes id_A) \circ \tilde{T}_2 \circ (id_A \otimes T) = T \circ (\mu \otimes id_A), \quad (4.2)$$

$$\tilde{T}_1 \circ (T \otimes id_A) \circ (id_A \otimes T) = \tilde{T}_2 \circ (id_A \otimes T) \circ (T \otimes id_A). \quad (4.3)$$

*Then $(A, \mu \circ T, u)$ is also an algebra in $\mathcal{C}$, denoted by A^T . The morphism T is called a **pseudotwistor** and the two morphisms $\tilde{T}_1, \tilde{T}_2$ are called the **companions** of T . If $\mathcal{C}$ is the category of k -vector spaces, $\tilde{T}_1 = \tilde{T}_2 = T_{13}$ and $T_{12} \circ T_{23} = T_{23} \circ T_{12}$, then T is called a **twistor** for A .*

Proposition 4.2 *Let $\mathcal{C}$ be a monoidal category and T a laycle on $\mathcal{C}$. If (A, μ, u) is an algebra in $\mathcal{C}$, then $T_{A,A}$ is a pseudotwistor for A , with companions $\tilde{T}_1 := T_{A,A,A}^b$ and $\tilde{T}_2 := T_{A,A,A}^f$, where T^b and T^f are the families defined by (2.3) and (2.4).*

Proof. We prove (4.1). The naturality of T implies $T_{A,A} \circ (id_A \otimes \mu) = (id_A \otimes \mu) \circ T_{A,A \otimes A}$, and from (2.3) we obtain $T_{A,A} \circ (id_A \otimes \mu) = (id_A \otimes \mu) \circ T_{A,A,A}^b \circ (T_{A,A} \otimes id_A)$, q.e.d. Similarly one can prove (4.2), while (4.3) follows immediately by using (2.3), (2.4) and (2.2). $\square$

Corollary 4.3 *If T is a laycle on a monoidal category $\mathcal{C}$ and (A, μ, u) is an algebra in $\mathcal{C}$, then $(A, \mu \circ T_{A,A}, u)$ is also an algebra in $\mathcal{C}$.*

Remark 4.4 *If $\mathcal{C}$ is a monoidal category and c is a braiding on $\mathcal{C}$, then, by [5], the double braiding $c_{X,Y}^2 := c_{Y,X} \circ c_{X,Y}$ is a twine on $\mathcal{C}$, in particular a laycle. Thus, Proposition 4.2 generalizes the fact (proved in [24], Corollary 6.8) that a double braiding induces a pseudotwistor on every algebra in $\mathcal{C}$.*

Definition 4.5 *Let $\mathcal{C}$ be a monoidal category, (A, μ, u) an algebra in $\mathcal{C}$ and $T : A \otimes A \rightarrow A \otimes A$ a pseudotwistor with companions $\tilde{T}_1$ and $\tilde{T}_2$. We say that T is a **strong pseudotwistor** if T is invertible and the following conditions are satisfied:*

$$\tilde{T}_2 \circ (id_A \otimes T) = (id_A \otimes T) \circ \tilde{T}_1, \quad (4.4)$$

$$\tilde{T}_1 \circ (T \otimes id_A) = (T \otimes id_A) \circ \tilde{T}_2. \quad (4.5)$$

In this case, we denote

$$T_{A \otimes A, A} := \tilde{T}_2 \circ (id_A \otimes T) = (id_A \otimes T) \circ \tilde{T}_1,$$

$$T_{A, A \otimes A} := \tilde{T}_1 \circ (T \otimes id_A) = (T \otimes id_A) \circ \tilde{T}_2.$$

Remark 4.6 *If $T_{X,Y}$ is a laycle on a monoidal category $\mathcal{C}$ and (A, μ, u) is an algebra in $\mathcal{C}$, then, by (2.2), it follows that $T_{A,A}$ is a strong pseudotwistor for A .*

Lemma 4.7 *If T is a strong pseudotwistor, then the following relations hold:*

$$(T \otimes id_A) \circ T_{A \otimes A, A} = T_{A \otimes A, A} \circ (T \otimes id_A), \quad (4.6)$$

$$(id_A \otimes T) \circ T_{A, A \otimes A} = T_{A, A \otimes A} \circ (id_A \otimes T), \quad (4.7)$$

$$T_{A \otimes A, A} \circ (T \otimes id_A) = T_{A, A \otimes A} \circ (id_A \otimes T), \quad (4.8)$$

$$(T \otimes id_A) \circ \tilde{T}_2 \circ (id_A \otimes T) = (id_A \otimes T) \circ \tilde{T}_1 \circ (T \otimes id_A). \quad (4.9)$$

Proof. Straightforward computation, using (4.4), (4.5) and (4.3). $\square$

Our next results are the analogues for pseudotwistors of the facts that composition of laycles is a laycle and the inverse of a laycle is a laycle.

Proposition 4.8 *Let $\mathcal{C}$ be a monoidal category, (A, μ, u) an algebra in $\mathcal{C}$ and $T, D : A \otimes A \rightarrow A \otimes A$ two strong pseudotwistors for A , such that*

$$D_{A, A \otimes A} \circ (id_A \otimes T) = (id_A \otimes T) \circ D_{A, A \otimes A}, \quad (4.10)$$

$$D_{A \otimes A, A} \circ (T \otimes id_A) = (T \otimes id_A) \circ D_{A \otimes A, A}. \quad (4.11)$$

Then $U := T \circ D$ is a pseudotwistor for A , with companions $\tilde{U}_1 := T_{A, A \otimes A} \circ \tilde{D}_1 \circ (T^{-1} \otimes id_A)$ and $\tilde{U}_2 := T_{A \otimes A, A} \circ \tilde{D}_2 \circ (id_A \otimes T^{-1})$. If moreover we have

$$T_{A, A \otimes A} \circ (id_A \otimes D) = (id_A \otimes D) \circ T_{A, A \otimes A}, \quad (4.12)$$

$$T_{A \otimes A, A} \circ (D \otimes id_A) = (D \otimes id_A) \circ T_{A \otimes A, A}, \quad (4.13)$$

then U is also a strong pseudotwistor.

Proof. We check (4.1)–(4.3) for U :

$$\begin{aligned}
U \circ (id_A \otimes \mu) &= T \circ D \circ (id_A \otimes \mu) \\
&\stackrel{(4.1)}{=} (id_A \otimes \mu) \circ \tilde{T}_1 \circ (T \otimes id_A) \circ \tilde{D}_1 \circ (D \otimes id_A) \\
&= (id_A \otimes \mu) \circ T_{A,A \otimes A} \circ \tilde{D}_1 \circ (D \otimes id_A) \\
&= (id_A \otimes \mu) \circ \tilde{U}_1 \circ (U \otimes id_A),
\end{aligned}$$

$$\begin{aligned}
U \circ (\mu \otimes id_A) &= T \circ D \circ (\mu \otimes id_A) \\
&\stackrel{(4.2)}{=} (\mu \otimes id_A) \circ \tilde{T}_2 \circ (id_A \otimes T) \circ \tilde{D}_2 \circ (id_A \otimes D) \\
&= (\mu \otimes id_A) \circ T_{A \otimes A, A} \circ \tilde{D}_2 \circ (id_A \otimes D) \\
&= (\mu \otimes id_A) \circ \tilde{U}_2 \circ (id_A \otimes U),
\end{aligned}$$

$$\begin{aligned}
\tilde{U}_1 \circ (U \otimes id_A) \circ (id_A \otimes U) &= T_{A,A \otimes A} \circ \tilde{D}_1 \circ (D \otimes id_A) \circ (id_A \otimes T) \circ (id_A \otimes D) \\
&= T_{A,A \otimes A} \circ D_{A,A \otimes A} \circ (id_A \otimes T) \circ (id_A \otimes D) \\
&\stackrel{(4.10)}{=} T_{A,A \otimes A} \circ (id_A \otimes T) \circ D_{A,A \otimes A} \circ (id_A \otimes D) \\
&\stackrel{(4.8)}{=} T_{A \otimes A, A} \circ (T \otimes id_A) \circ D_{A \otimes A, A} \circ (D \otimes id_A) \\
&\stackrel{(4.11)}{=} T_{A \otimes A, A} \circ D_{A \otimes A, A} \circ (T \otimes id_A) \circ (D \otimes id_A) \\
&= T_{A \otimes A, A} \circ \tilde{D}_2 \circ (id_A \otimes D) \circ (T \otimes id_A) \circ (D \otimes id_A) \\
&= \tilde{U}_2 \circ (id_A \otimes U) \circ (U \otimes id_A),
\end{aligned}$$

proving that U is a pseudotwistor for A . We assume now that (4.12) and (4.13) hold and we prove (4.4) and (4.5) for U :

$$\begin{aligned}
(id_A \otimes U) \circ \tilde{U}_1 &= (id_A \otimes T) \circ (id_A \otimes D) \circ T_{A,A \otimes A} \circ \tilde{D}_1 \circ (T^{-1} \otimes id_A) \\
&\stackrel{(4.12)}{=} (id_A \otimes T) \circ T_{A,A \otimes A} \circ (id_A \otimes D) \circ \tilde{D}_1 \circ (T^{-1} \otimes id_A) \\
&= (id_A \otimes T) \circ T_{A,A \otimes A} \circ D_{A \otimes A, A} \circ (T^{-1} \otimes id_A) \\
&\stackrel{(4.11)}{=} (id_A \otimes T) \circ T_{A,A \otimes A} \circ (T^{-1} \otimes id_A) \circ D_{A \otimes A, A} \\
&\stackrel{(4.8), (4.7)}{=} T_{A \otimes A, A} \circ D_{A \otimes A, A} \\
&= T_{A \otimes A, A} \circ \tilde{D}_2 \circ (id_A \otimes D) \\
&= \tilde{U}_2 \circ (id_A \otimes U),
\end{aligned}$$

$$\begin{aligned}
(U \otimes id_A) \circ \tilde{U}_2 &= (T \otimes id_A) \circ (D \otimes id_A) \circ T_{A \otimes A, A} \circ \tilde{D}_2 \circ (id_A \otimes T^{-1}) \\
&\stackrel{(4.13)}{=} (T \otimes id_A) \circ T_{A \otimes A, A} \circ (D \otimes id_A) \circ \tilde{D}_2 \circ (id_A \otimes T^{-1}) \\
&= (T \otimes id_A) \circ T_{A \otimes A, A} \circ D_{A,A \otimes A} \circ (id_A \otimes T^{-1}) \\
&\stackrel{(4.10)}{=} (T \otimes id_A) \circ T_{A \otimes A, A} \circ (id_A \otimes T^{-1}) \circ D_{A,A \otimes A} \\
&\stackrel{(4.8), (4.6)}{=} T_{A,A \otimes A} \circ D_{A,A \otimes A} \\
&= T_{A,A \otimes A} \circ \tilde{D}_1 \circ (D \otimes id_A) \\
&= \tilde{U}_1 \circ (U \otimes id_A),
\end{aligned}$$

showing that U is a strong pseudotwistor. $\square$

Corollary 4.9 *If T is a strong pseudotwistor for an algebra (A, μ, u) in a monoidal category $\mathcal{C}$, then $T \circ T$ is also a strong pseudotwistor for A .*

Proposition 4.10 *Let $\mathcal{C}$ be a monoidal category, (A, μ, u) an algebra in $\mathcal{C}$ and $T : A \otimes A \rightarrow A \otimes A$ a strong pseudotwistor for A such that the companions $\tilde{T}_1$ and $\tilde{T}_2$ are invertible. Then the inverse $V := T^{-1}$ is also a strong pseudotwistor for A , with companions $\tilde{V}_1 = \tilde{T}_2^{-1}$ and $\tilde{V}_2 = \tilde{T}_1^{-1}$.*

Proof. Straightforward computation, using (4.1)–(4.3) for T together with (4.4) and (4.5). $\square$

Remark 4.11 *Let $\mathcal{C}$ be a monoidal category, (A, μ, u) an algebra in $\mathcal{C}$, $T : A \otimes A \rightarrow A \otimes A$ a strong pseudotwistor for A and $\mathcal{D}$ a laycle on $\mathcal{C}$. Then T and $D := \mathcal{D}_{A,A}$ satisfy (4.10) and (4.11), hence $T \circ D$ is a pseudotwistor for A .*

Our next result is the analogue for pseudotwistors of the fact that if σ, σ' are cohomologous lazy cocycles on a Hopf algebra H then the algebras $H(\sigma)$ and $H(\sigma')$ are isomorphic:

Proposition 4.12 *Let $\mathcal{C}$ be a monoidal category, (A, μ, u) an algebra in $\mathcal{C}$, $T : A \otimes A \rightarrow A \otimes A$ a strong pseudotwistor for A and $R_X : X \rightarrow X$ a family of natural isomorphisms in $\mathcal{C}$ such that $R_I = id_I$. Then we have an algebra isomorphism*

$$R_A : A^{T \circ D^1(R)_{A,A}} \simeq A^T.$$

Proof. Note first that $T \circ D^1(R)_{A,A}$ is a pseudotwistor by Remark 4.11. We compute:

$$\begin{aligned} R_A \circ \mu \circ T \circ D^1(R)_{A,A} &= R_A \circ \mu \circ T \circ R_{A \otimes A}^{-1} \circ (R_A \otimes R_A) \\ &\stackrel{\text{naturalness of } T}{=} \mu \circ R_{A \otimes A} \circ T \circ R_{A \otimes A}^{-1} \circ (R_A \otimes R_A) \\ &\stackrel{\text{naturalness of } R}{=} \mu \circ T \circ (R_A \otimes R_A), \end{aligned}$$

finishing the proof. $\square$

If T is a laycle on a monoidal category $\mathcal{C}$, then, by [5], T is a twine if and only if the following condition is satisfied:

$$(T_{X,Y,Z}^f \otimes id_W) \circ (id_X \otimes T_{Y,Z,W}^b) = (id_X \otimes T_{Y,Z,W}^b) \circ (T_{X,Y,Z}^f \otimes id_W), \quad (4.14)$$

for all $X, Y, Z, W \in \mathcal{C}$. Note that the families T^f, T^b coincide respectively to the families A, B from the Definition 1.5 of a pure-braided structure, and (4.14) coincides with (1.13). We are thus led to the following concept and terminology:

Definition 4.13 *Let $\mathcal{C}$ be a monoidal category, A an algebra in $\mathcal{C}$ and $T : A \otimes A \rightarrow A \otimes A$ a pseudotwistor with companions $\tilde{T}_1$ and $\tilde{T}_2$. We call T a **pure pseudotwistor** if*

$$(\tilde{T}_2 \otimes id_A) \circ (id_A \otimes \tilde{T}_1) = (id_A \otimes \tilde{T}_1) \circ (\tilde{T}_2 \otimes id_A). \quad (4.15)$$

Corollary 4.14 *A twine on a monoidal category $\mathcal{C}$ induces a pure pseudotwistor on every algebra A in $\mathcal{C}$. In particular, if c is a braiding on $\mathcal{C}$ then $c_{A,A}^2$ is a pure pseudotwistor for A .*

Remark 4.15 Obviously, a pseudotwistor for which $\tilde{T}_1 = \tilde{T}_2 = id_{A \otimes A \otimes A}$ is pure. Here are some concrete (but nonunital) examples of such pseudotwistors:

- (i) take A an associative algebra, $R = R^1 \otimes R^2 \in A \otimes A$ and define $T(a \otimes b) = aR^1 \otimes R^2b$, for all $a, b \in A$.
- (ii) take A an associative algebra, $f : A \rightarrow A$ a linear map satisfying $f(ab) = af(b)$ for all $a, b \in A$, and $T(a \otimes b) = f(a) \otimes b$. If instead f satisfies $f(ab) = f(a)b$, then take $T(a \otimes b) = a \otimes f(b)$.
- (iii) take A an associative algebra, $\delta : A \rightarrow A \otimes A$ a linear map such that $\delta(ab) = (a \otimes 1)\delta(b)$ for all $a, b \in A$ and $T : A \otimes A \rightarrow A \otimes A$, $T(a \otimes b) = \delta(a)(1 \otimes b)$. If instead δ satisfies $\delta(ab) = \delta(a)(1 \otimes b)$, then take $T(a \otimes b) = (a \otimes 1)\delta(b)$.

Note that example (i) was inspired by a construction in [2], while (ii) and (iii) are related to some constructions in [22] involving so-called (anti-) dipterous algebras.

Example 4.16 If A is an associative algebra and $T : A \otimes A \rightarrow A \otimes A$ is a twistor, then it is easy to see that T is pure.

Example 4.17 We recall some facts from [24]. Let (Ω, d) be a DG algebra, that is $\Omega = \bigoplus_{n \geq 0} \Omega^n$ is a graded algebra and $d : \Omega \rightarrow \Omega$ is a linear map with $d(\Omega^n) \subseteq \Omega^{n+1}$ for all $n \geq 0$, $d^2 = 0$ and $d(\omega\zeta) = d(\omega)\zeta + (-1)^{|\omega|}\omega d(\zeta)$ for all homogeneous ω and ζ , where $|\omega|$ is the degree of ω . The Fedosov product ([16], [11]), given by $\omega \circ \zeta = \omega\zeta - (-1)^{|\omega|}d(\omega)d(\zeta)$, for homogeneous ω and ζ , gives a new associative algebra structure on Ω . We consider $\mathcal{C}$ to be the monoidal category of $\mathbb{Z}_2$ -graded vector spaces, and regard Ω as a $\mathbb{Z}_2$ -graded algebra (i.e. an algebra in $\mathcal{C}$) by putting even components in degree zero and odd components in degree one. Define the linear map

$$T : \Omega \otimes \Omega \rightarrow \Omega \otimes \Omega, \quad T(\omega \otimes \zeta) = \omega \otimes \zeta - (-1)^{|\omega|}d(\omega) \otimes d(\zeta),$$

for homogeneous ω and ζ . Then T is a pseudotwistor for Ω in $\mathcal{C}$, affording the Fedosov product. Its companions are given (for homogeneous ω, ζ, η) by

$$\tilde{T}_1(\omega \otimes \zeta \otimes \eta) = \tilde{T}_2(\omega \otimes \zeta \otimes \eta) = \omega \otimes \zeta \otimes \eta - (-1)^{|\omega|+|\zeta|}d(\omega) \otimes \zeta \otimes d(\eta).$$

We claim that T is a pure pseudotwistor. Indeed, a straightforward computation shows that

$$\begin{aligned} (\tilde{T}_2 \otimes id) \circ (id \otimes \tilde{T}_1)(\omega \otimes \zeta \otimes \eta \otimes \nu) &= (id \otimes \tilde{T}_1) \circ (\tilde{T}_2 \otimes id)(\omega \otimes \zeta \otimes \eta \otimes \nu) \\ &= \omega \otimes \zeta \otimes \eta \otimes \nu - (-1)^{|\omega|+|\zeta|}d(\omega) \otimes \zeta \otimes d(\eta) \otimes \nu \\ &\quad - (-1)^{|\zeta|+|\eta|}\omega \otimes d(\zeta) \otimes \eta \otimes d(\nu) - (-1)^{|\omega|+|\eta|}d(\omega) \otimes d(\zeta) \otimes d(\eta) \otimes d(\nu), \end{aligned}$$

for all homogeneous ω, ζ, η, ν .

We recall the following result from [24]:

Proposition 4.18 ([24]) *Let (A, μ, u) be an algebra in a monoidal category $\mathcal{C}$, let $R, P : A \otimes A \rightarrow A \otimes A$ twisting maps between A and itself such that R is invertible, and assume that*

$$(P \otimes id_A) \circ (id_A \otimes P) \circ (P \otimes id_A) = (id_A \otimes P) \circ (P \otimes id_A) \circ (id_A \otimes P), \quad (4.16)$$

$$(R \otimes id_A) \circ (id_A \otimes R) \circ (R \otimes id_A) = (id_A \otimes R) \circ (R \otimes id_A) \circ (id_A \otimes R), \quad (4.17)$$

$$(P \otimes id_A) \circ (id_A \otimes P) \circ (R \otimes id_A) = (id_A \otimes R) \circ (P \otimes id_A) \circ (id_A \otimes P), \quad (4.18)$$

$$(R \otimes id_A) \circ (id_A \otimes P) \circ (P \otimes id_A) = (id_A \otimes P) \circ (P \otimes id_A) \circ (id_A \otimes R). \quad (4.19)$$

Define $T : A \otimes A \rightarrow A \otimes A$, $T := R^{-1} \circ P$. Then T is a pseudotwistor with companions

$$\tilde{T}_1 = (R^{-1} \otimes id_A) \circ (id_A \otimes T) \circ (R \otimes id_A), \quad \tilde{T}_2 = (id_A \otimes R^{-1}) \circ (T \otimes id_A) \circ (id_A \otimes R).$$

Our next result is the analogue for pseudotwistors of the fact from [5] that a family of the type $T_{X,Y} = c'_{Y,X} \circ c_{X,Y}$, with c, c' braidings, is a twine:

Proposition 4.19 *Assume that the hypotheses of Proposition 4.18 hold. Then:*

- (i) *T is a pure pseudotwistor;*
- (ii) *assume that moreover P is also invertible and*

$$(P \otimes id_A) \circ (id_A \otimes R) \circ (R \otimes id_A) = (id_A \otimes R) \circ (R \otimes id_A) \circ (id_A \otimes P), \quad (4.20)$$

$$(R \otimes id_A) \circ (id_A \otimes R) \circ (P \otimes id_A) = (id_A \otimes P) \circ (R \otimes id_A) \circ (id_A \otimes R) \quad (4.21)$$

(these conditions appear in [24] too and they imply that R is also a twisting map between A^T and itself). Then T is a strong pseudotwistor.

Proof. We check (4.15):

$$\begin{aligned} (\tilde{T}_2 \otimes id_A) \circ (id_A \otimes \tilde{T}_1) &= (id_A \otimes R^{-1} \otimes id_A) \circ (T \otimes id_A \otimes id_A) \circ (id_A \otimes R \otimes id_A) \\ &\quad \circ (id_A \otimes R^{-1} \otimes id_A) \circ (id_A \otimes id_A \otimes T) \circ (id_A \otimes R \otimes id_A) \\ &= (id_A \otimes R^{-1} \otimes id_A) \circ (id_A \otimes id_A \otimes T) \circ (T \otimes id_A \otimes id_A) \\ &\quad \circ (id_A \otimes R \otimes id_A) \\ &= (id_A \otimes R^{-1} \otimes id_A) \circ (id_A \otimes id_A \otimes T) \circ (id_A \otimes R \otimes id_A) \\ &\quad \circ (id_A \otimes R^{-1} \otimes id_A) \circ (T \otimes id_A \otimes id_A) \circ (id_A \otimes R \otimes id_A) \\ &= (id_A \otimes \tilde{T}_1) \circ (\tilde{T}_2 \otimes id_A). \end{aligned}$$

Assume now that P is invertible and (4.20), (4.21) hold. Obviously T is invertible, and we only have to check (4.4) and (4.5):

$$\begin{aligned} \tilde{T}_2 \circ (id_A \otimes T) &= (id_A \otimes R^{-1}) \circ (T \otimes id_A) \circ (id_A \otimes R) \circ (id_A \otimes T) \\ &= (id_A \otimes R^{-1}) \circ (R^{-1} \otimes id_A) \circ (P \otimes id_A) \circ (id_A \otimes P) \\ &\stackrel{(4.20)}{=} (id_A \otimes R^{-1}) \circ (id_A \otimes P) \circ (R^{-1} \otimes id_A) \circ (id_A \otimes R^{-1}) \\ &\quad \circ (P^{-1} \otimes id_A) \circ (id_A \otimes R) \circ (P \otimes id_A) \circ (id_A \otimes P) \\ &\stackrel{(4.18)}{=} (id_A \otimes R^{-1}) \circ (id_A \otimes P) \circ (R^{-1} \otimes id_A) \\ &\quad \circ (id_A \otimes R^{-1}) \circ (id_A \otimes P) \circ (R \otimes id_A) \\ &= (id_A \otimes T) \circ \tilde{T}_1, \end{aligned}$$

$$\begin{aligned} \tilde{T}_1 \circ (T \otimes id_A) &= (R^{-1} \otimes id_A) \circ (id_A \otimes T) \circ (R \otimes id_A) \circ (T \otimes id_A) \\ &= (R^{-1} \otimes id_A) \circ (id_A \otimes R^{-1}) \circ (id_A \otimes P) \circ (P \otimes id_A) \\ &\stackrel{(4.21)}{=} (R^{-1} \otimes id_A) \circ (P \otimes id_A) \circ (id_A \otimes R^{-1}) \circ (R^{-1} \otimes id_A) \\ &\quad \circ (id_A \otimes P^{-1}) \circ (R \otimes id_A) \circ (id_A \otimes P) \circ (P \otimes id_A) \\ &\stackrel{(4.19)}{=} (R^{-1} \otimes id_A) \circ (P \otimes id_A) \circ (id_A \otimes R^{-1}) \\ &\quad \circ (R^{-1} \otimes id_A) \circ (P \otimes id_A) \circ (id_A \otimes R) \\ &= (T \otimes id_A) \circ \tilde{T}_2, \end{aligned}$$

finishing the proof. □

Remark 4.20 If T is a braided twistor as introduced in [24], a computation identical to the one in the proof of Proposition 4.19 (i) shows that T is a pure pseudotwistor.

Example 4.21 Let A be an algebra and $\mathcal{F}$ a braid system over A as introduced by Durdevich in [14], that is a collection of bijective twisting maps between A and itself, satisfying the condition

$$(\alpha \otimes id_A) \circ (id_A \otimes \beta) \circ (\gamma \otimes id_A) = (id_A \otimes \gamma) \circ (\beta \otimes id_A) \circ (id_A \otimes \alpha), \quad \forall \alpha, \beta, \gamma \in \mathcal{F}.$$

For $\alpha, \beta \in \mathcal{F}$ define the map $T_{\alpha, \beta} : A \otimes A \rightarrow A \otimes A$, $T_{\alpha, \beta} := \alpha^{-1} \circ \beta$. By [24] we know that T is a pseudotwistor for A , and by Proposition 4.19 it follows that it is a pure strong pseudotwistor.

We introduce now the categorical version of Borchers' R -matrices:

Proposition 4.22 Let $\mathcal{C}$ be a monoidal category, (A, μ, u) an algebra in $\mathcal{C}$ and $T : A \otimes A \rightarrow A \otimes A$ a morphism in $\mathcal{C}$ such that $T \circ (u \otimes id_A) = u \otimes id_A$ and $T \circ (id_A \otimes u) = id_A \otimes u$. Assume that there exist two morphisms $\bar{T}_1, \bar{T}_2 : A \otimes A \otimes A \rightarrow A \otimes A \otimes A$ in $\mathcal{C}$ such that

$$(id_A \otimes \mu) \circ (T \otimes id_A) \circ \bar{T}_1 = T \circ (id_A \otimes \mu), \quad (4.22)$$

$$(\mu \otimes id_A) \circ (id_A \otimes T) \circ \bar{T}_2 = T \circ (\mu \otimes id_A), \quad (4.23)$$

$$(T \otimes id_A) \circ \bar{T}_1 \circ (id_A \otimes T) = (id_A \otimes T) \circ \bar{T}_2 \circ (T \otimes id_A). \quad (4.24)$$

Then $(A, \mu \circ T, u)$ is also an algebra in $\mathcal{C}$, denoted by A^T . The morphism T is called an **R -matrix** and the two morphisms $\bar{T}_1, \bar{T}_2$ are called the **companions** of T . Obviously, the original concept of R -matrix is obtained for $\mathcal{C}$ being the category of k -vector spaces and $\bar{T}_1 = \bar{T}_2 = T_{13}$.

Proof. Obviously u is a unit for $(A, \mu \circ T)$; we check the associativity of $\mu \circ T$:

$$\begin{aligned} (\mu \circ T) \circ ((\mu \circ T) \otimes id_A) &= (\mu \circ T) \circ (\mu \otimes id_A) \circ (T \otimes id_A) \\ &\stackrel{(4.23)}{=} \mu \circ (\mu \otimes id_A) \circ (id_A \otimes T) \circ \bar{T}_2 \circ (T \otimes id_A) \\ &\stackrel{(4.24)}{=} \mu \circ (\mu \otimes id_A) \circ (T \otimes id_A) \circ \bar{T}_1 \circ (id_A \otimes T) \\ &= \mu \circ (id_A \otimes \mu) \circ (T \otimes id_A) \circ \bar{T}_1 \circ (id_A \otimes T) \\ &\stackrel{(4.22)}{=} \mu \circ T \circ (id_A \otimes \mu) \circ (id_A \otimes T) \\ &= (\mu \circ T) \circ (id_A \otimes (\mu \circ T)), \end{aligned}$$

finishing the proof. $\square$

Proposition 4.23 Let $\mathcal{C}$ be a monoidal category, (A, μ, u) an algebra in $\mathcal{C}$ and $T : A \otimes A \rightarrow A \otimes A$ an invertible morphism in $\mathcal{C}$. Then T is a pseudotwistor if and only if it is an R -matrix. More precisely, if T is a pseudotwistor with companions $\tilde{T}_1, \tilde{T}_2$ then T is an R -matrix with companions $\bar{T}_1 = (T^{-1} \otimes id_A) \circ \tilde{T}_1 \circ (T \otimes id_A)$ and $\bar{T}_2 = (id_A \otimes T^{-1}) \circ \tilde{T}_2 \circ (id_A \otimes T)$; conversely, if T is an R -matrix with companions $\bar{T}_1, \bar{T}_2$ then T is a pseudotwistor with companions $\tilde{T}_1 = (T \otimes id_A) \circ \bar{T}_1 \circ (T^{-1} \otimes id_A)$ and $\tilde{T}_2 = (id_A \otimes T) \circ \bar{T}_2 \circ (id_A \otimes T^{-1})$.

Proof. Straightforward computation. $\square$

Corollary 4.24 Let $\mathcal{C}$ be a monoidal category and T a laycle on $\mathcal{C}$. If (A, μ, u) is an algebra in $\mathcal{C}$, then $T_{A,A}$ is an R -matrix for A , with companions $\bar{T}_1 := T_{A,A,A}^f$ and $\bar{T}_2 := T_{A,A,A}^b$, where T^b and T^f are the families defined by (2.3) and (2.4).

5 A characterization of generalized double braidings

Let $\mathcal{C}$ be a monoidal category and A an algebra in $\mathcal{C}$. If T is a pseudotwistor for A and $R : A \otimes A \rightarrow A \otimes A$ is an invertible twisting map such that the companions of T are given by the formulae

$$\tilde{T}_1 = (R^{-1} \otimes id_A) \circ (id_A \otimes T) \circ (R \otimes id_A), \quad (5.1)$$

$$\tilde{T}_2 = (id_A \otimes R^{-1}) \circ (T \otimes id_A) \circ (id_A \otimes R), \quad (5.2)$$

then, by [24], Theorem 6.6, it follows that $R \circ T$ is a twisting map between A and itself. This result has the following categorical analogue, with laycles replacing pseudotwistors and braidings replacing twisting maps:

Theorem 5.1 *Let $\mathcal{C}$ be a monoidal category, T a laycle and d a braiding on $\mathcal{C}$, such that for all $X, Y, Z \in \mathcal{C}$ the following relations hold:*

$$T_{X \otimes Y, Z} = (id_X \otimes T_{Y, Z}) \circ (d_{X, Y}^{-1} \otimes id_Z) \circ (id_Y \otimes T_{X, Z}) \circ (d_{X, Y} \otimes id_Z), \quad (5.3)$$

$$T_{X, Y \otimes Z} = (T_{X, Y} \otimes id_Z) \circ (id_X \otimes d_{Y, Z}^{-1}) \circ (T_{X, Z} \otimes id_Y) \circ (id_X \otimes d_{Y, Z}). \quad (5.4)$$

Then the families $d'_{X, Y} := d_{X, Y} \circ T_{X, Y}$ and $d''_{X, Y} := T_{Y, X} \circ d_{X, Y}$ are also braidings on $\mathcal{C}$. Moreover, T is a twine and $d''_{X, Y} = T_{Y, X} \circ d'_{X, Y} \circ T_{X, Y}^{-1}$ (thus $(\mathcal{C}, d')$ and $(\mathcal{C}, d'')$ are braided isomorphic).

Proof. Note first that (5.3) and (5.4) are the analogues of (5.1) and (5.2), because they are respectively equivalent to

$$T_{X, Y, Z}^b = (d_{X, Y}^{-1} \otimes id_Z) \circ (id_Y \otimes T_{X, Z}) \circ (d_{X, Y} \otimes id_Z),$$

$$T_{X, Y, Z}^f = (id_X \otimes d_{Y, Z}^{-1}) \circ (T_{X, Z} \otimes id_Y) \circ (id_X \otimes d_{Y, Z}).$$

Also, as consequences of (2.2), (5.3) and (5.4) we obtain the following relations:

$$T_{X, Y \otimes Z} = (d_{X, Y}^{-1} \otimes id_Z) \circ (id_Y \otimes T_{X, Z}) \circ (d_{X, Y} \otimes id_Z) \circ (T_{X, Y} \otimes id_Z), \quad (5.5)$$

$$T_{X \otimes Y, Z} = (id_X \otimes d_{Y, Z}^{-1}) \circ (T_{X, Z} \otimes id_Y) \circ (id_X \otimes d_{Y, Z}) \circ (id_X \otimes T_{Y, Z}). \quad (5.6)$$

Now we check (1.6) and (1.7) for d' :

$$\begin{aligned} d'_{X, Y \otimes Z} &= d_{X, Y \otimes Z} \circ T_{X, Y \otimes Z} \\ &\stackrel{(1.6), (5.4)}{=} (id_Y \otimes d_{X, Z}) \circ (d_{X, Y} \otimes id_Z) \circ (T_{X, Y} \otimes id_Z) \circ (id_X \otimes d_{Y, Z}^{-1}) \\ &\quad \circ (T_{X, Z} \otimes id_Y) \circ (id_X \otimes d_{Y, Z}) \\ &\stackrel{(5.5)}{=} (id_Y \otimes d_{X, Z}) \circ (id_Y \otimes T_{X, Z}) \circ (d_{X, Y} \otimes id_Z) \circ (T_{X, Y} \otimes id_Z) \\ &= (id_Y \otimes d'_{X, Z}) \circ (d'_{X, Y} \otimes id_Z), \end{aligned}$$

$$\begin{aligned} d'_{X \otimes Y, Z} &= d_{X \otimes Y, Z} \circ T_{X \otimes Y, Z} \\ &\stackrel{(1.7), (5.3)}{=} (d_{X, Z} \otimes id_Y) \circ (id_X \otimes d_{Y, Z}) \circ (id_X \otimes T_{Y, Z}) \circ (d_{X, Y}^{-1} \otimes id_Z) \\ &\quad \circ (id_Y \otimes T_{X, Z}) \circ (d_{X, Y} \otimes id_Z) \\ &\stackrel{(5.6)}{=} (d_{X, Z} \otimes id_Y) \circ (T_{X, Z} \otimes id_Y) \circ (id_X \otimes d_{Y, Z}) \circ (id_X \otimes T_{Y, Z}) \\ &= (d'_{X, Z} \otimes id_Y) \circ (id_X \otimes d'_{Y, Z}). \end{aligned}$$

Thus, d' is a braiding. It is obvious that $d''_{X,Y} = T_{Y,X} \circ d'_{X,Y} \circ T_{X,Y}^{-1}$, and it follows that d'' is also a braiding, by using Proposition 2.8. The fact that T satisfies (1.17) follows immediately by using (5.5) and (5.6). $\square$

Corollary 5.2 *Let H be a Hopf algebra, $R \in H \otimes H$ a quasitriangular structure and $F \in H \otimes H$ a lazy twist, such that*

$$(\Delta \otimes id)(F) = F_{23} R_{12}^{-1} F_{13} R_{12}, \quad (5.7)$$

$$(id \otimes \Delta)(F) = F_{12} R_{23}^{-1} F_{13} R_{23}. \quad (5.8)$$

Then the elements $R' = RF$ and $R'' = F_{21}R$ are also quasitriangular structures on H .

Proposition 5.3 *Let $\mathcal{C}$ be a monoidal category and c, c' braidings on $\mathcal{C}$. Then the inverse braiding $d_{X,Y} := c_{Y,X}^{-1}$ and the laycle $T_{X,Y} = c'_{Y,X} \circ c_{X,Y}$ satisfy the hypotheses of Theorem 5.1. Consequently, the family $d'_{X,Y} = d_{X,Y} \circ T_{X,Y} = c_{Y,X}^{-1} \circ c'_{Y,X} \circ c_{X,Y}$ is a braiding on $\mathcal{C}$, and the braiding d'' coincides with the original braiding c' .*

Proof. We check (5.3):

$$\begin{aligned} & (id_X \otimes T_{Y,Z}) \circ (d_{X,Y}^{-1} \otimes id_Z) \circ (id_Y \otimes T_{X,Z}) \circ (d_{X,Y} \otimes id_Z) \\ &= (id_X \otimes c'_{Z,Y}) \circ (id_X \otimes c_{Y,Z}) \circ (c_{Y,X} \otimes id_Z) \circ (id_Y \otimes c'_{Z,X}) \\ & \quad \circ (id_Y \otimes c_{X,Z}) \circ (c_{Y,X}^{-1} \otimes id_Z) \\ & \stackrel{(1.6), \text{ naturality of } c}{=} (id_X \otimes c'_{Z,Y}) \circ (c'_{Z,X} \otimes id_Y) \circ (id_Z \otimes c_{Y,X}) \circ (c_{Y,Z} \otimes id_X) \\ & \quad \circ (id_Y \otimes c_{X,Z}) \circ (c_{Y,X}^{-1} \otimes id_Z) \\ & \stackrel{(1.8)}{=} (id_X \otimes c'_{Z,Y}) \circ (c'_{Z,X} \otimes id_Y) \circ (c_{X,Z} \otimes id_Y) \circ (id_X \otimes c_{Y,Z}) \\ & \quad \circ (c_{Y,X} \otimes id_Z) \circ (c_{Y,X}^{-1} \otimes id_Z) \\ & \stackrel{(1.6), (1.7)}{=} c'_{Z,X \otimes Y} \circ c_{X \otimes Y, Z} \\ &= T_{X \otimes Y, Z}. \end{aligned}$$

The proof of (5.4) is similar and left to the reader. $\square$

Remark 5.4 *Theorem 5.1 together with Proposition 5.3 provide an alternative proof of the fact from [5] that the laycle $T_{X,Y} = c'_{Y,X} \circ c_{X,Y}$ is a twine.*

Remark 5.5 *If $(\mathcal{C}, c)$ is a braided monoidal category and we take the inverse braiding $d_{X,Y} = c_{Y,X}^{-1}$, then in general $(\mathcal{C}, c)$ and $(\mathcal{C}, d)$ are not braided isomorphic. Thus, the braidings d' and d'' obtained in Theorem 5.1 are in general not equivalent to the original braiding d .*

Theorem 5.1 together with Proposition 5.3 provide the following characterization of generalized double braidings:

Proposition 5.6 *Let $\mathcal{C}$ be a monoidal category and T a laycle on $\mathcal{C}$. Then T is a generalized double braiding if and only if there exists a braiding d on $\mathcal{C}$ such that (5.3) and (5.4) hold.*

References

- [1] J. Bichon, G. Carnovale, Lazy cohomology: an analogue of the Schur multiplier for arbitrary Hopf algebras, *J. Pure Appl. Algebra* **204** (2006), 627–665.
- [2] J. Bichon, R. Street, Militaru’s D -equation in monoidal categories, *Appl. Categ. Structures* **11** (2003), 337–357.
- [3] R. E. Borcherds, Vertex algebras, In *Topological field theory, primitive forms and related topics*, pp. 35–77, Progr. Math. **160**, Birkhauser, Boston, 1998.
- [4] R. E. Borcherds, Quantum vertex algebras, In *Adv. Stud. Pure Math.* **31**, pp. 51–74, Math. Soc. Japan, Tokyo, 2001.
- [5] A. Bruguières, Double braidings, twists and tangle invariants, *J. Pure Appl. Algebra* **204** (2006), 170–194.
- [6] G. Carnovale, Some isomorphisms for the Brauer groups of a Hopf algebra, *Comm. Algebra* **29** (2001), 5291–5305.
- [7] G. Carnovale. The Brauer group of modified supergroup algebras, *J. Algebra* **305** (2006), 993–1036.
- [8] G. Carnovale, J. Cuadra, Cocycle twisting of $E(n)$ -module algebras and applications to the Brauer group, *K-Theory* **33** (2004), 251–276.
- [9] H. X. Chen, Skew pairing, cocycle deformations and double crossproducts, *Acta Math. Sinica, English Ser.* **15** (1999), 225–234.
- [10] J. Cuadra, F. Panaite, Extending lazy 2-cocycles on Hopf algebras and lifting projective representations afforded by them, *J. Algebra* **313** (2007), 695–723.
- [11] J. Cuntz, D. Quillen, Algebra extensions and nonsingularity, *J. Amer. Math. Soc.* **8** (1995), 251–289.
- [12] V. G. Drinfeld, Quantum groups, In *Proc. Int. Cong. Math. (Berkeley, 1986)*, pp. 798–820, Amer. Math. Soc., Providence, RI, 1987.
- [13] V. G. Drinfeld, Quasi-Hopf algebras, *Leningrad Math. J.* **1** (1990), 1419–1457.
- [14] M. Durdevich, Generalized braided quantum groups, *Israel J. Math.* **98** (1997), 329–348.
- [15] P. Etingof, S. Gelaki, Quasisymmetric and unipotent tensor categories, preprint arXiv:math.QA/07081487.
- [16] B. V. Fedosov, Analytical formulas for the index of an elliptic operator, *Trans. Moscow Math. Soc.* **30** (1974), 159–240.
- [17] S. Gelaki, On pointed ribbon Hopf algebras, *J. Algebra* **181** (1996), 760–786.
- [18] L. M. Ionescu, Cohomology of monoidal categories and non-abelian group cohomology, PhD thesis, 2000, available on <http://www.ilstu.edu>.

- [19] P. Jara Martínez, J. López Peña, F. Panaite, F. Van Oystaeyen, On iterated twisted tensor products of algebras, preprint arXiv:math.QA/0511280, to appear in *Internat. J. Math.*
- [20] A. Joyal, R. Street, Braided tensor categories, *Adv. Math.* **102** (1993), 20–78.
- [21] C. Kassel, "Quantum groups", *Graduate Texts in Mathematics* **155**, Springer Verlag, Berlin, 1995.
- [22] P. Leroux, On some remarkable operads constructed from Baxter operators, preprint arXiv:math.QA/0311214.
- [23] G. Liu, S. Zhu, Almost-triangular Hopf algebras, *Algebr. Represent. Theory* **10** (2007), 555–564.
- [24] J. López Peña, F. Panaite, F. Van Oystaeyen, General twisting of algebras, *Adv. Math.* **212** (2007), 315–337.
- [25] S. Majid, Anyonic quantum groups, In *Spinors, twistors, Clifford algebras and quantum deformations*, pp. 327–336, Kluwer Academic Publishers, 1993.
- [26] S. Majid, "Foundations of quantum group theory", Cambridge Univ. Press, 1995.
- [27] F. Panaite, F. Van Oystaeyen, Quasitriangular structures for some pointed Hopf algebras of dimension 2^n , *Comm. Algebra* **27** (1999), 4929–4942.
- [28] F. Panaite, F. Van Oystaeyen, Clifford-type algebras as cleft extensions for some pointed Hopf algebras, *Comm. Algebra* **28** (2000), 585–600.
- [29] F. Panaite, M. D. Staic, F. Van Oystaeyen, On some classes of lazy cocycles and categorical structures, *J. Pure Appl. Algebra* **209** (2007), 687–701.
- [30] B. Pareigis, Symmetric Yetter-Drinfeld categories are trivial, *J. Pure Appl. Algebra* **155** (2001), 91.
- [31] D. E. Radford, On the antipode of a quasitriangular Hopf algebra, *J. Algebra* **151** (1992), 1–11.
- [32] D. E. Radford, Minimal quasitriangular Hopf algebras, *J. Algebra* **157** (1993), 285–315.
- [33] P. Schauenburg, Hopf bimodules, coquasibialgebras, and an exact sequence of Kac, *Adv. Math.* **165** (2002), 194–263.
- [34] M. D. Staic, Pure-braided Hopf algebras and knot invariants, *J. Knot Theory Ramifications* **13** (2004), 385–400.
- [35] F. Van Oystaeyen, Y. Zhang, The Brauer group of a braided monoidal category, *J. Algebra* **202** (1998), 96–128.